



AVERTISSEMENT

Ce document est le fruit d'un long travail approuvé par le jury de soutenance et mis à disposition de l'ensemble de la communauté universitaire élargie.

Il est soumis à la propriété intellectuelle de l'auteur. Ceci implique une obligation de citation et de référencement lors de l'utilisation de ce document.

D'autre part, toute contrefaçon, plagiat, reproduction illicite encourt une poursuite pénale.

Contact : ddoc-theses-contact@univ-lorraine.fr

LIENS

Code de la Propriété Intellectuelle. articles L 122. 4

Code de la Propriété Intellectuelle. articles L 335.2- L 335.10

http://www.cfcopies.com/V2/leg/leg_droi.php

<http://www.culture.gouv.fr/culture/infos-pratiques/droits/protection.htm>



UNIVERSITÉ
DE LORRAINE

UFR mathématique informatique
mécanique et automatique

UNIVERSITY OF NAIROBI



CBPS
School of Mathematics

Ecole Doctorale IAEM – Lorraine

Mathematical Analysis and Dynamical Systems : Modeling of Highland Malaria in Western Kenya.

A Joint Dissertation Submitted In Partial Fulfillment of the
Requirements For the Degree of DOCTOR OF PHILOSOPHY
In THE UNIVERSITY OF LORRAINE and THE UNIVERSITY OF NAIROBI
Specialization : Applied Mathematics

Defense Date :23/11/2012

By

Josephine W. KAGUNDA

Review Committee Members :

1. ARINO Julien, Professor – University of Manitoba
2. AUGER Pierre, Professor – IRD Bondy ED P VI
3. GOUZE Jean-Luc, HDR - INRIA, Sophia Antipolis

Examiners :

1. SALLET Gauthier, Professor - University of Lorraine-Supervisor
2. OGANA Wandera, Professor - University of Nairobi- Supervisor

TABLE DES MATIÈRES

Appreciation	v
Dedication	viii
Abstract	ix
Resume	1
Introduction	2
0.0.1 Highland Malaria	2
0.0.2 Mathematical Modeling	8
1 Some Epidemiological Models	11
1.1 The First Model : D.Bernoulli	11
1.1.1 The Model	12
1.1.2 Analysis of the model	13
1.2 An SIR Deterministic Model :Kermack and Mc Kendrick	14
1.2.1 Model Formulation	14
1.2.2 Analysis of the model	15
1.3 Transmission of DHF in an Age Structured Population	16
1.4 Differential Susceptibility and Infectivity by Hyman	19
1.5 A model for Disease transmission in a Metapopulation by J.Arino	20
1.6 Ross Model	22
1.7 Global analysis of a vector-host epidemic model with nonlinear incidence by Li-Ming Cai et. al.	23
1.8 Ross-Macdonald Model in a Patchy environment by Auger et. al.	25
2 A vector SIS malaria model in a patchy environment with age structure	27
Introduction	27
2.1 Introduction	27
2.2 The model	29

2.3	Reduced System	31
2.4	Basic properties of the model	36
3	Stability Analysis	40
3.1	Basic Reproduction Number, $\mathcal{R}_0$	40
3.2	Main Result	41
3.3	An Example in Two Patches	45
3.4	Simulation	46
3.4.1	Parameter values	47
3.5	Numerical simulation	47
3.5.1	The U-shaped Valley Sites : Iguhu and Umutete, $\mathcal{R}_0 = 5.36$	48
3.5.2	The V-shaped Valley Sites : Fort Tenan and Marani, $\mathcal{R}_0 = 1.67$	51
3.6	Discussion and Biological implications	51
3.7	Conclusion	52
4	Dynamics of Vector-Host feeding Contact Rate with Saturation in malaria	54
4.1	Introduction	54
4.2	Vector-Host Contact with saturation	56
4.3	The Model Equations	58
5	Stability Analysis : Saturation	61
5.1	Global stability of the Disease Free Equilibrium	61
5.1.1	A compact positively invariant set	61
5.1.2	Reduction of the model	62
5.1.3	Basic reproduction ratio	62
5.1.4	Equilibria	63
5.1.5	Global Stability of the DFE	64
5.1.6	Stability of the endemic equilibrium	65
5.1.7	Global stability of the endemic equilibrium, when there is no disease induced mortality	68
5.1.8	Simulation	68
5.1.9	Estimating Q_v	68
5.2	Discussion and Conclusion	69
6	Conclusion and Perspective	73
A	Mathematical Tools	75
A.1	Preliminaries and Notation	75
A.2	Stability	76
A.2.1	Stability : Basic definition	76
A.2.2	Dynamical Properties	77
A.3	Monotone systems	79
A.3.1	Monotone Dynamical System	79
A.3.2	Triangular System	80
A.3.3	Lyapunov Methods	82
A.3.4	LaSalle Invariance Principle	83
A.4	Matrices	83
A.4.1	Metzler Matrices	86

A.4.2	Characteristics of Metzler stable matrices	89
A.4.3	Irreducible Metzler Matrices	90
A.4.4	Regular Splitting of a Metzler matrix	92
A.5	Calculation of Basic Reproduction Number	93
B	Biology of mosquito and malaria parasite	95
B.0.1	The parasites life cycle.	95
B.0.2	Mosquitoes Life cycle	95

TABLE DES FIGURES

1	The Distribution of Malaria in Kenya (MARA/ARMA)	3
2	The malaria endemic Western Kenya Highlands. The areas affected by epidemics are 1500-2500 meters above sea level [50]	4
3	A picture of an U-shaped Valley [161]	6
4	A picture of a V-shaped Valley [161]	7
5	The antibodies prevalence for the U-shaped and the V-shaped valleys [161]	7
2.1	Transfer Diagram for the different epidemiological groups in different patches . .	39
3.1	A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the U-shaped valleys system. The age structure in the populations is clearly shown. In this case $\mathcal{R}_0 = 5.36$	49
3.2	A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the U-shaped valleys ecosystem. A case where there is no age structure in the population. In this case $\mathcal{R}_0 = 5.36$	49
3.3	A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the U-shaped valleys ecosystem. The is no age structure and the two ecosystems are treated as one U-shaped valley ecosystem	50
3.4	A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the V-shaped valleys ecosystem with age structure. In this case $\mathcal{R}_0 = 0.96 < 1$	50
3.5	A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the V-shaped valleys ecosystem. The is no age structures in the populations. In this case $\mathcal{R}_0 = 0.96 < 1$	51
4.1	A graphical comparison between f_1 and f_2	57
5.1	Domain of study	63
5.2	Variation of susceptible host and vector populations at the Disease Free equilibrium. $Q_v = 0.3$ and $\tilde{\mathcal{R}}_0 = 0.718 < 1$	71
5.3	Variation of susceptible host and vector populations at the Endemic equilibrium. $Q_v = 0.09$ and $\tilde{\mathcal{R}}_0 = 2.495 > 1$	71
5.4	Variation of Infected Host, Infected Vector, Total Host and Total Vector populations at the Disease Free equilibrium. $Q_v = 0.3$ and $\tilde{\mathcal{R}}_0 = 0.718 < 1$	72

5.5	Variation of Infected Host, Infected Vector, Total Host and Total Vector populations at the Endemic Equilibrium. $Q_v = 0.09$ and $\hat{\mathcal{R}}_0 = 2.495 > 1$	72
A.1	Vector fields pointing toward the interior	78

Appreciation

"What can I do, since the words of my mouth are not enough, to tell you the things you have done for me, for they are wonderful. My heart will praise you Oh God, I will not forget your benefits."

This doctoral thesis has been successful courtesy, of the financial and institutional support offered by the French Embassy in Nairobi and IRD, the University of Nairobi and the Laboratory of Mathematics and Applications at the University of Lorraine, within the project team MASAIE, INRIA. The supervision was carried out jointly between the two universities. I would like to thank all those who have been directly or indirectly involved towards this noble course and achievement.

There are many people that I would like to thank for making this dissertation possible. First of all, I thank my two Supervisors, Prof. SALLET Gauthier and Prof. OGANA Wandera, who created the many ideas and directions for the research, suggested methods for the analysis, verified the results, and proofread my numerous drafts. I thank them for all their advice, both, mathematical and non-mathematical ; for the time they have given to this research ; for giving me a chance to work with them ; and most of all, for making the process of writing this dissertation enjoyable. To Sallet and Ogana, 'You modeled the mathematician in me, you are an excellent example of integrity and patriotism that I desire to follow. It is your remarks that have enabled me to improve the quality of this work'.

I wish to thank Dr. Were and Dr. Nyandwi who were very instrumental in putting me in this Joint Thesis program on behalf of the University of Nairobi. Very often they spent their precious time advising, encouraging and facilitating me during my research. I am very grateful to Prof. IGGDR Abderrahman, who often offered his support in the LMAM during my research in Metz. Many thanks to Prof. Marion Mutugi who introduced me to the world of malaria in 2009. I am so grateful to Mme. Christel Wiemert for the dedicated support she accorded me at INRIA. She made me feel at home in MASAIE and always provided the necessary tools I needed to work. My appreciation also goes to Service de l'Information Scientifique et Technique (IST) who went out of their way to provided me with books and papers that I required for the research. I thank all members of the School of Mathematics, University of Nairobi who never missed an encouraging word whenever we met. Many are the moments we held fruitful discussions that kept me going especially when the going got tough.

Many thanks to my Final Oral Defense Examination Committee for carefully reading the dissertation and for their suggestions on improving it. I also thank my reviewers for reading the dissertation. Special thanks to Dr. Githeko of KEMRI for the many malaria insights and data that he gave us during the writing of this thesis.

I take this opportunity to thank all the MASAIE Team comrades, with whom I had a pleasant stay in Metz : Bichara, Leontine, Lena, Plaire and Berge. The generosity of these members our my team has allowed me to borrow priceless knowledge A lot of thanks to my friends and colleagues : James Okwoyo, Dossou, Aime, and members of L'Eglise Evangelique du Metz for allowing me to spend my sundays as they should be spent.

I thank my children, my brothers and sister, who prayed, encouraged me and wanted to see this thesis become a reality, suffering for long during my absence. I thank especially of my Dad, Allan

who at his age made it his duty to do all he could to make my studies a success. Last but not the least, I wish to thank Nancy, who became a mother to my children while I was away and never got tired of meeting their maternal needs, as she kept assuring me that all is well so that I concentrate, even when one of them would be seriously ill and on medication, it is only God who can pay her for her services.

Dedication

I dedicate this Thesis to my children :

- Elijah Mbugua
- Favour Lilly Wamuca and
- Victor Mwaniki

For the suffering you endured during my long and repeated absence. May you find consolation and comfort enough to ease the grief it has caused you.

Abstract

The objective of this thesis is to model highland malaria in Western Kenya using dynamical systems. Two mathematical models are formulated; one, on differentiated susceptibility and differentiated infectivity in a metapopulation setting with age structure, the other, a saturated vector feeding rate model with disease induced deaths and varying host and vector populations. In the first model, we consider the different ecosystems identified as malaria hotspots in the Western Kenya highlands and consider the ecosystems as different patches. The population in each patch is classified as, either child (age 0-5 years) or, adult (over 5 years of age). Our results are compared with some published results in [161].

The model will aid in examining the role of ecosystem heterogeneity and age structure to the persistent malaria epidemics in the highlands. We formulate the differentiated susceptibility and infectivity model that extend to multiple patches the well known epidemiological models in one patch.

Classifying the hot spots as n patches, we give its mathematical analysis using the theory of triangular system, monotone non-linear dynamical systems, and Lyapunov-Lasalle invariance principle techniques. Key to our analysis is the definition of a reproductive number, $\mathcal{R}_0$, the number of new infections caused by one individual in an otherwise fully susceptible population throughout the duration of the infectious period. The existence and stability of disease-free and endemic equilibrium is established. We prove that the disease free state of the systems is globally asymptotically stable when the basic reproduction number $\mathcal{R}_0 \leq 1$, and when $\mathcal{R}_0 > 1$ an endemic equilibrium is established which is locally and globally asymptotically stable. The model shows that the age structuring reduces the magnitude of infection. Using relevant data we did some simulation, to demonstrate the role played by metapopulation and age structuring on the incidence and $\mathcal{R}_0$.

In the second part we formulate a model for malaria with saturation on the vector feeding rates that lead to a nonlinear function in the infection term. The vector feeding rate is assumed, as in the predator prey models, to rise linearly as a function of the host-vector ratio until it reaches a threshold Q_v , after which the vector feeds freely at its desired rate. The two populations (host and vector) are variable and drive malaria transmission, such that when the vectors are fewer than hosts, the rate of feeding is determined by the vectors feeding desire, whereas, when the hosts are more than the vectors, the feeding rate is limited by host availability and other feeding sources may have to be sought by the vector. Malaria induced deaths are introduced in the host population, while the vector is assumed to survive with the parasite till its death. We prove that the Disease Free Equilibrium is locally and globally asymptotically stable if $\mathcal{R}_0 \leq 1$ and when $\mathcal{R}_0 > 1$, an endemic equilibrium emerges, which is unique, locally and globally asymptotically stable. The role of the saturated mosquito feeding rate is explored with simulation showing the crucial role it plays especially on the basic reproduction number.

Key Words : Modeling, Metapopulation, Differentiated infectivity and Susceptibility, Monotone Dynamical Systems, Lyapunov methods, Basic Reproduction Number, Global stability, Saturation, Malaria, Numerical Simulation.

Resume

L'objectif de cette thèse est de modéliser la transmission du paludisme dans la région montagneuse de l'Ouest du Kenya, en se servant des outils de systèmes dynamiques. Nous considérons deux modèles mathématiques. Le premier prend en compte une susceptibilité et une infectivité différentielle dans les métapopulations, et le second un taux de saturation des repas sanguins dans la population des moustiques.

Dans le premier modèle, nous considérons plusieurs écosystèmes identifiés comme zones sensibles dans la région montagneuse de l'Ouest du Kenya. Dans ce modèle, ces zones sensibles sont considérées comme nos différents patchs. Les populations de chaque patch sont divisées en deux : les enfants (0 à 5 ans), et les adultes (dont l'âge est supérieur à 5 ans). Les résultats obtenus sont comparés avec ceux obtenus dans [161]. Le modèle nous permet d'évaluer le rôle de l'hétérogénéité de l'écosystème et la persistance de l'épidémie dans la région, due à la structuration d'âge. Nous prenons en compte la susceptibilité et l'infectivité différentielle afin d'étendre le modèle d'un patch en un modèle à plusieurs patchs.

Après avoir subdivisé la région en n zones sensibles, nous faisons une analyse mathématique du modèle obtenu. Pour effectuer cette analyse, nous utilisons la théorie des systèmes triangulaires, des systèmes dynamiques monotones, des systèmes dynamiques non linéaires anti-monotones et le principe d'invariance de LaSalle. Un des éléments très utilisés dans notre analyse qui est un concept clé en épidémiologie, est le taux de reproduction de base, très souvent noté $\mathcal{R}_0$. Cette quantité, sans dimension, est le nombre moyen de cas secondaires, engendré par un individu infectieux typique durant sa période d'infectiosité, quand il est introduit dans une population constituée entièrement de susceptibles. L'existence et la stabilité du point d'équilibre sans maladie (DFE) sont établies et nous prouvons que le DFE est globalement asymptotiquement stable lorsque $\mathcal{R}_0 \leq 1$. Lorsque $\mathcal{R}_0 > 1$, le modèle admet un point d'équilibre endémique qui est globalement asymptotiquement stable. L'analyse de notre modèle montre que la structuration d'âge réduit l'ampleur de l'infection. En utilisant les données relevées, nous faisons quelques simulations numériques afin montrer l'impact de la métapopulation et de la structuration d'âge sur le taux de reproduction de base.

Dans la seconde partie, nous formulons un modèle de paludisme avec saturation du taux d'alimentation des moustiques qui nous conduit à une incidence non linéaire. Nous démontrons que le DFE est globalement asymptotiquement stable si $\mathcal{R}_0 < 1$. Lorsque $\mathcal{R}_0 > 1$, il existe un unique point d'équilibre endémique qui est globalement asymptotiquement stable. Des simulations numériques sont faites afin d'illustrer l'impact de la saturation du taux d'alimentation sur le taux de reproduction de base.

Mots clés : Modélisation, métapopulation, susceptibilité et infectivité différentielle, systèmes dynamiques monotones, méthode de Lyapunov, taux de reproduction de base, stabilité globale, saturation, paludisme, simulations numériques.

INTRODUCTION

0.0.1 Highland Malaria

Malaria is a protozoan disease caused by a parasite transmitted by a female anopheles mosquito, as it seeks a blood meal for development of its egg. The disease is common in the East African region, Kenya included. Malaria is one of the leading causes of morbidity and mortality in Kenya and it kills an estimated 34,000 children under five in Kenya every year. In Kenya 77% of the population lives in areas where malaria is transmitted. Of all the visits in Kenyan hospitals, malaria is responsible for 30% of all outpatient visits (requiring more than eight million out-patient treatments at health facilities each year) and 15% of all hospital admissions. About 3.5 million children are at risk of infection and developing severe malaria. Pregnant women also face high risks. There are approximately 1.1 million pregnancies per year in malaria endemic areas. During pregnancy, malaria can cause miscarriage and anemia. Each year, an estimated 6,000 pregnant women suffer from malaria-associated anemia, and 4,000 babies are born with low birth weight as a result of maternal anemia. Economically, it is estimated that 170 million working days in Kenya are lost each year because of malaria illness.

Malaria did not exist in the Western Kenya highlands until the second decade of the 20th century, so the highlands were regarded as safe havens from the surrounding malarious areas of Uganda and Kenya [110, 79]. After the first World War, malaria was prevalent in the communities inhabiting the highlands. This was as a result of wide-scale population settlement resulting from the completion of the railway line from Kenyan coast via the highlands to Lake Victoria. This also facilitated the gradual spread of infective mosquitoes into the highlands from the low-lying hyperendemic-disease areas [45]. The development of tea estates and agriculture in the highlands, with the concomitant clearing of the forests, provided suitable mosquito breeding grounds. Finally, importation of infected labourers completed the conditions necessary for malaria transmission [107].

This resulted to frequent malaria epidemics, with some authors labeling the resurgent epidemics as 'Highland malaria', because they viewed it as a new typology variant, demanding special attention in the new global commitment to Roll Back Malaria [165]. 'Highland' malaria therefore, either is a new phenomenon or a reemergence of a previous prevailing epidemiology [136].

Highland malaria in Western Kenya is characterized by unstable and high transmission varia-




Figure 1 – The Distribution of Malaria in Kenya (MARA/ARMA)

bility which results in epidemics during periods of suitable climatic conditions. The sensitivity of a site to malaria epidemics depends on the level of immunity of the human population. The epidemics in the highlands have continued to wreck havoc on the public health of the inhabitants, resulting in high morbidity and mortality especially in children under five years [136]. Since the early 1980s, there have been massive percentage increases in *P. falciparum* burden at African highland locations [58]. The epidemics varies seasonally and spatially with many studies being carried out to ascertain the drivers of such variations. Among the factors considered as key drivers of the resurgent epidemics that characterise the highlands include but are not limited to the following :

1. Climate Variation.

Climate variation plays an important role in initiating malaria epidemics in the Western Kenya highlands. The developmental rates and survivorship of the malaria parasite and the




Figure 2 – The malaria endemic Western Kenya Highlands. The areas affected by epidemics are 1500-2500 meters above sea level [50]

malaria vector is highly temperature dependent, while rainfall influences the availability of breeding places, hence their demography. Small changes in temperature could therefore provide transiently suitable conditions for unstable transmission within populations that have acquired little functional immunity. Studies on climatic variations and malaria epidemics in Western Kenya show a high spatial variation in the sensitivity to malaria outpatient number to climate fluctuations in the highlands [169]. The contribution of climate variation to reemergence of malaria epidemics has been controversial, and a good conclusion is that other factors together with climate change are more plausible explanations for the malaria resurgence.

2. Drug Resistance

During the past two decades, Chloroquine (CQ) resistance has spread geographically, increased in prevalence within its range, and intensified in its severity of clinical failure [144, 164]. CQ resistance has been identified as an important factor in the malaria resurgence on the Kericho tea estates [46]. How much effect this one has on the magnitude of the epidemics is a point of discussion; this is because drug resistance can only aggravate malaria induced morbidity and mortality not initiating an epidemic. Again, drug resistance cannot explain the sporadic malaria epidemics in the Kenyan highlands in the 1920s and

1950s, when the problem of drug resistance was insignificant [49].

3. Land Use Changes.

The increasing human populations in the East African highlands put a lot of pressure on the sensitive ecosystem [104]. There is an increase in changing human activities such as deforestation and swamp cultivation for subsistence agriculture, growing cash crops, and firewood acquisition that have characterized the region.

Reports indicate that there is an 8% reduction in forest cover in the East African highlands in one decade [1]. The changes in land use, especially deforestation in the highlands, have significantly changed the microclimatic conditions of mosquito breeding sites that have significantly had effects on the life history of the vectors.

Deforestation is said to have enhanced vectorial capacity by 10.6% and 29% in the dry and rainy seasons in the highlands, respectively, thereby resulting in high indoor temperatures. This causes faster blood meal digestion, thereby shortening the length of the gonotrophic cycle of mosquitoes. Reduced gonotrophic cycle length would cause mosquitoes to feed more often and lay eggs more frequently in a deforested area. [1].

4. Population Growth and Health Service Provision

An important factor to consider in the escalating malaria incidence in Western Kenya is population growth. Of vital consideration is the relationship between the percentage population growth rate in the districts served by each health facility and the percentage rises in malaria cases. Over the last 3 decades, sub-Saharan Africa including the highlands, has seen a high rate of increase in population size, resulting from high fertility rates and increasing child survival with annual growth rates averaging 3.9%. This implies that, holding the incidence of malaria constant, the disease would be expected to have doubled over approximately an 18-year period. Clearly, without a concomitant investment in essential clinical services, beds, staff, and supporting infrastructure, the changing requirements for clinical management will have been perceived by most district-level public health officials as a crisis [57].

Defining true epidemics is difficult [136]. For most public health workers, epidemics represent exacerbations of disease out of proportion to the normal level to which that facility is subject; these increases overwhelm the facilities ability to cope. Therefore, a slow but pervasive epidemic of clinical malaria may have emerged in the highlands, where lack of investment in the physical capacity to manage an increasing population has resulted inevitably in more malaria cases that require a basic clinical service.

5. Immunity.

Due to endemicity of malaria in Africa, people are infected so frequently that after a long time, they develop a degree of acquired immunity, making them asymptomatic carriers of infection [13]. The people who use mosquito nets in the night are not generally infected by the disease since the small number of bites they receive when they are outside the nets are not sufficient to cause the disease, but rather, the small exposure to infection gives them immunity. The low level exposure to infection, however, is important as it acts as vaccination and develops immunity against the disease [86]. Those who acquire the immunity

are however susceptible to reinfections because the immune protection may wane over time (temporary immunity) or may not be fully protective (partial immunity) [146].

Studies in Western Kenya indicate that in areas of unstable transmission in the highlands, the prevalence of circumsporozoite protein (CSP) is about 13% in adults over 40 years of age whereas in the stable transmission lowlands, approximately 65% of children are antibody positive [2]. This shows that the human population in the highlands has fewer people with immunity and this renders them vulnerable to severe forms of malaria during epidemics [161]. Since the risk of an epidemic is closely related to the level of immunity of the human population, the final outcome of the epidemic will be closely linked to the proportion of people who may not have had exposure to the disease and who have no immunity to malaria.

6. Ecosystem heterogeneity.

A study [161] on the different ecosystems in the Western Kenya highlands identify three major characteristics : the plateaus, V-shaped valleys and U-shaped valleys. The V-shaped valleys have steep hillsides and narrow valleys that allow fast flow of the rivers, while the U-shaped valleys have flat bottomed valleys with slow moving rivers.

! "#\$%& ' (") % * +



! "#\$%&' (") % * +



Figure 4 – A picture of a V-shaped Valley [161]

transmission of the parasites.




Figure 5 – The antibodies prevalence for the U-shaped and the V-shaped valleys [161]

The plateau’s antibody and parasite prevalence was similar to that of the V-shaped valleys. Although climatic variability drives malaria epidemics in Western Kenya highlands the terrain characteristics can modify the level of malaria transmission and the rate of immunity development.

It is our aim in this thesis to formulate a mathematical model that captures the heterogeneity in the terrain characteristics and the infection dynamics of this very important disease in the economy of Kenya.

0.0.2 Mathematical Modeling

Mathematical models have become important tools in analyzing the spread and control of infectious diseases. The model formulation process clarifies assumptions, variables, and parameters; moreover, models provide conceptual results such as thresholds, basic reproduction numbers, contact numbers, and replacement numbers. Understanding the transmission characteristics of infectious diseases in communities, regions, and countries can lead to better approaches to decreasing the transmission of these diseases.

Mathematical modeling arises from the idea that spread of infectious disease and transmission dynamics follow laws that can be formulated in mathematical language.

In a mathematical model the populations are subdivided into homogenous epidemiological groups called compartments. Compartmental labels such as M, S, E, I, and R are often used for the epidemiological classes. Depending on the nature of the disease, we can have different compartments for the susceptibles, infected and infectious, recovered or immune individuals and so on. An SIS disease is one in which susceptibles (S) become infected (I) and recover without immunity and so are again susceptible an SIR disease in which a susceptible (S) becomes infected (I) and infecteds that recover are immune (R). If the immune individual spends some time in the recovered compartment to return again to the susceptible compartment, we have a SIRS model. If there is some latent period between the time of infection before becoming infectious, the model becomes SEIRS. Other types of include MSEIR, MSEIRS, SEIR, SEIRS, SIR, SIRS, SEI, SEIS, SI. In this thesis, we concentrate on SIS models for the sake of malaria. Its true that no permanent immunity is achieved for malaria and even the partial or acquired immunity does not protect infection, it only reduces the severity of the disease to the infectee [63].

Mathematical modeling begun with Daniel Bernoulli in 1766 when he modeled the effects of smallpox variolation (a precursor of vaccination) on life expectancy [31]. In the beginning of that century there was much discussion about why an epidemic ended before all susceptible were infected with hypotheses about changing virulence of the pathogen during the epidemic. Sir Ronald Ross, who received the Nobel prize in 1902 for elucidating the life cycle of the malaria parasite, used mathematical modeling to investigate the effectiveness of various intervention strategies for malaria [128]. Kermack and McKendrick described the dynamics of disease transmission in terms of a system of differential equations in [83, 84, 85]. They were the first to use the so-called basic reproduction number, a threshold that determines if a disease dies out or spreads in a susceptible population. Literature review in [17, 34, 37, 61] shows the rapid growth of epidemiology modeling towards the end of the twentieth century as mathematical modeling became more popular to aid in public health policy making.

Many mathematical models describing the dynamics of AIDS pandemic have been used to predict the course of the epidemic and identify the most effective prevention strategies. The need for evaluating intervention strategies for newly emerging and re-emerging pathogens has revealed the real impact of mathematical modeling on public health.

There are two types of mathematical models, deterministic and stochastic. The deterministic models are transmission models using ordinary differential equations to describe the process, whereas stochastic models use statistical mechanisms to describe the transmission of infection between two individuals. The rigorous mathematical analyses of the processes contribute to the

understanding of disease transmission in much detail. Therefore, developing a mathematical model helps address the essential processes involved in shaping the epidemiology of an infectious disease and reveals the parameters that are most important and crucial for control and data collection. Mathematical modeling is then also integrative in combining knowledge from different disciplines like microbiology, social sciences, and clinical sciences [91].

Of special interest in this thesis is metapopulation models, which refer to many small and extinction-prone local populations connected via migration in fragmented environment. Metapopulation theory is based on Levin's multi-groups model [102]. The population is subdivided into a finite number of discrete subpopulations or metapopulations called patches that are assumed to be homogeneous and well mixed. Then each patch in the population is subdivided into compartments corresponding to different epidemiological status. This leads to multi-patch, multi-compartmental systems. There are three possible formulations. The first assumes that an individual in a patch of infection can infect a susceptible individual in another patch. This approach assumes that there is a short distance between patches, but individuals (human or mosquito) do not migrate between patches (they can make brief visits sufficient for transmission). This assumption leads to a series of models that has been studied in [63, 72, 96]. The legendary model of Lajmanovich and Yorke [96] is a key example of such models.

The second formulation considers the mobility of individuals between patches with two forms of movement : Migration, that is change of respective patch residence. The situation is that of a digraph or directed movement graph. For the short visits, the individual visit a number of patches but returns to the individual patch of residence. The "Mover -Stayer " movement is covered in the mode of Sattenspiel and Dietz [133]. Here, the formulation takes care of the degree of patches connectivity. Recently there has been an increasing interest in this type of models [7, 9, 11, 29, 44, 80, 159, 160].

The third formulation is more recent and takes into account the network structure of the metapopulations, that is those patches that (local populations) are locally connected. This formulation takes into account the degree of connectivity (number of links between a patch and its immediate neighbours) of a patch and models the mobility of individuals in terms of diffusion between discrete patches. This is a statistical approach where the system states depends on the local population densities of the metapopulation network. This leads to a situation of a non-oriented graph connectivity. [130, 131, 154, 155, 156].

The rest of the thesis is arranged as follows.

In the first part, we construct and analyze mathematically an age structured model for highland malaria in Western Kenya, with differentiated patch susceptibility and infectivity and human migration between patches. We show that there is a threshold below which the disease vanishes and above which the disease is persistent in the metapopulation. It is applied to two sites in Western Kenya that have been identified as 'malaria hotspots' in literature cited above. This part contributes significantly to mathematics since little has been done on the rigorous analysis of these types of models or metapopulation.

In the second part, we formulate a model for the dynamics of vector feeding habits with saturation and varying host and vector population. We establish mathematically the existence of a disease free equilibrium where the disease dies out and an endemic equilibrium, where the disease

persists in the population using $\mathcal{R}_0$ as a threshold. The local and endemic equilibria are proved to be locally and globally asymptotically stable. The analysis of the role of saturation in biting rate is also done to show that it plays a key role in infection and also incidence of malaria.

SOME EPIDEMIOLOGICAL MODELS

In this section we give a review of relevant literature that motivated the work done in this thesis. The selection is done from the first mathematical models and formulation of SIS models. Other interesting reviews have been done on age structured models, differentiated infectivity models, saturation and meta-population models. Finally we have looked Ross model for malaria and a model for malaria in a patchy environment. Throughout this section we will verify that these models are well-posed and they form a foundation for the models we formulate in this thesis. We begin with the model formulated by D. Bernoulli on smallpox.

1.1 The First Model : D.Bernoulli

The main objective of Bernoulli was to calculate the gain in life expectancy at birth if smallpox were to be eliminated as a cause of death. Smallpox is an endemic, potentially lethal, infectious disease that is dreaded and feared. One in every five adults will die if infected by smallpox, (nearly one in three patients). If it does not cause death, smallpox often leaves a trail of destroyed lives. Effective treatment for smallpox has remained elusive. In the eleventh century, the Chinese practiced variolation, which is a form of inoculation, where infectious material from smallpox cases was transferred into the skin of susceptibles with the intention of inducing lifelong immunity by a mild infection with a low risk of death. The result was, however risky, because there was a possible rise in mortality rate by 1 to 2%. In 1721 this method was introduced from Turkey into England by the wife of the English Ambassador to Constantinople, Lady Mary Wortley Montague.

Smallpox was fortunately eradicated by the first human voluntary control. Its official "disappearance" dates back to 1977 (when the last case of smallpox was reported). This was made possible by the combined efforts of smallpox vaccination and screening program led by WHO in 1967.

Bernoulli demonstrated before the Academy of Sciences that, despite the dangers it presented in the fight against smallpox, the generalization of variolation imported from the East, would increase the life expectancy from 26 years and 7 months to 29 years and 9 months. His calculations have passed the test of time, and he was persuaded that his model was good. We give the details of the model he formulated.

1.1.1 The Model

Bernoulli assumed the following

- The probability that an individual infected by smallpox for the first time dies is p and the individual survives is $1 - p$ and this probability is independent of the individual's age.
- q is the probability that an individual is infected in the year and it is independent of his age. (ie the probability that an individual is infected during a small time interval dx between age x and age $x + dx$ is $q.dx$)
- when an individual survives after being infected with smallpox, he is immune for the rest of his life.

Let $m(x)$ be the natural mortality at age x , then the probability that an individual dies in a small time interval dx between age x and age $x + dx$ is $m(x).dx$. Considering a group of P_0 individuals born in the same year we have :

- $S(x)$ the number of individuals who are still alive at age x without having been infected (and therefore are more likely to be)
- $R(x)$ the number of immune individuals who are still alive at age x .
- $P(x) = S(x) + R(x)$ the total number of individuals who are still alive at age x .

When births correspond to $x = 0$, $S(0) = P(0) = P_0$ while between age x and age $x + dx$ (dx is infinitely small), an uninfected individual has a probability $q.dx$ of catching smallpox and a probability $m(x).dx$ of dying from another cause. So the variation in the number of individuals who are not yet infected is $dS(x) = -qS(x).dx - m(x)S(x).dx$, then we have the differential equation,

$$\frac{dS}{dx} = -qS(x) - m(x)S(x). \quad (1.1)$$

During the same small time interval, the number of individuals who die of smallpox is $p q S(x) dx$ and the number of survivors is $(1 - p) q S(x) dx$. In addition, there are $m(x) R(x)$ already immune individuals who suffer natural death, leading to a second differential equation

$$\frac{dR}{dx} = q(1 - p)S(x) - m(x)R(x). \quad (1.2)$$

Summing the above two equations we obtain

$$\frac{dP}{dx} = -p q S(x) - m(x)P(x). \quad (1.3)$$

From equations 1.1 and 1.3, Bernoulli showed that the fraction of individuals at age x who are at still at risk of contracting smallpox is

$$\frac{S(x)}{P(x)} = \frac{1}{(1 - p)e^{qx} + p}. \quad (1.4)$$

1.1.2 Analysis of the model

Bernoulli eliminated $m(x)$ between the equations 1.1 and 1.3 by expressing $m(x)$ as

$$-m(x) = q + \frac{1}{S} \frac{dS}{dx} = p q \frac{S}{P} + \frac{1}{P} \frac{dP}{dx}$$

to get the following expression

$$\frac{1}{P} \frac{dP}{dx} - \frac{S}{P^2} \frac{dP}{dx} = -q \frac{S}{P} + p q \left(\frac{S}{P}\right)^2$$

where the right hand side is the derivative of $f(x) = \frac{S(x)}{P(x)}$, and it is the proportion of infected individuals are who survive to age x . Therefore

$$\frac{df}{dx} = -q f + p q f^2. \quad (1.5)$$

The solution of this type of equation has been done by Jacques Bernoulli.

Let $g(x) = \frac{1}{f(x)}$, we see that $dg/dx = dg - p q$ and that $g(0) = 1/f(0) = 1$. We then set $h(x) = g(x) - p$, so that $dh/dx = q h$. So $h(x) = h(0) e^{qx} = (1-p) e^{qx}$. Finally, $g(x) = (1-p) e^{qx} + p$ and $f(x) = 1/g(x)$.

To illustrate his theory, Bernoulli used the mortality table by Halley (1656 - 1742) built for the city of Breslau (1693). This table listed the number of survivors at the beginning of the year $x(x = 1, 2, \dots)$ of a cohort of 1238 individuals born in year 0. Bernoulli however, believed like most of his contemporaries, that the numbers given by Halley were for the survivors reaching age x , that is $P(x)$ in the model. Because of this little confusion (Halley's article was indeed not very explicit), Bernoulli replaced the numbers 1238 by 1300 to obtain a realistic mortality for the first year of life.

Parameter values chosen by Bernoulli were

- The probability of dying from smallpox $p = \frac{1}{8}$
- The annual probability of catching smallpox $q = \frac{1}{8}$. Bernoulli then chose q so that the total death toll of smallpox represented $\frac{1}{13}$ of all deaths, which corresponded to the proportion found in many cities in Europe.

With the formula 1.4 and data values of $P(x)$ from the table of Halley we can calculate the number of individuals $S(x)$ who are still alive at age x without being infected, then determine easily the number of individuals who had smallpox and survived to age x : $R(x) = P(x) - S(x)$. Finally, we determine for each age x , the number of deaths due to smallpox between age x and age $x + 1$. In theory, it should be given by the integral $p q \int_x^{x+1} S(t) dt$ but the formula $p q (S(x) + S(x + 1))/2$ is a good approximation (Trapezium Rule). In total, Bernoulli deduced that out of the 1300 newborns, 101 were destined to die of smallpox, which is $1/13$ as he had calculated.

Bernoulli then considered a harmless way in which the whole population would be inoculated from small pox at birth. Smallpox would be eradicated and the question is, what the gain would be in life expectancy. Thus, starting from the same number P_0 at birth and noting $P^*(x)$ the number of individuals surviving at age x in the absence of smallpox, we have,

$$\frac{dP^*}{dx} = -m(x) P^* \quad (1.6)$$

Using equation 1.3, Bernoulli showed that

$$P^*(x) = \frac{P(x)}{1 - p + p P^* e^{-qx}}. \quad (1.7)$$

In particular the $P(x)/P^*(x) \rightarrow 1 - p$ for large values of x . One way to compare $P(x)$ and $P^*(x)$ is to estimate the life expectancy at birth, whose theoretical expression is

$$EP = \frac{1}{P_0} \int_0^\infty P(x) dx$$

with smallpox, and a similar expression with $P^*(x)$ instead of $P(x)$ in the absence of smallpox. Bernoulli used the approximate expression $(P(0) + P(1) + P(2) + \dots)/P_0$ which is similar to the one given by the trapezium rule. He finally found that life expectancy E with smallpox is 26 years and 7 months. Without smallpox, the life expectancy E^* is 29 years and 8 months.

Inoculation at birth would add more than 3 years to life expectancy. Of course, inoculation with a little virulent strain of smallpox, is not completely without risk. If p' is the probability of dying from smallpox during inoculation ($p' < p$), while life expectancy in the case where all children are inoculated at birth becomes $(1 - p')E^*$. This remains greater than the natural life expectancy E , if $p < 1 - E/E^*$, about 11%. Although there was no precise data available at the time, Bernoulli estimated the risk p , to be in fact less than 1%.

In the next section we look at the model by Kermack and Mc Kendrick whose theory is a mathematical hypothesis about how infectious diseases spread through a population. It is indeed the source of SIR models and their relatives.

1.2 An SIR Deterministic Model :Kermack and Mc Kendrick

1.2.1 Model Formulation

W.O. Kermack and AG Mc Kendrick [84, 83, 85] formulated a simple model of directly transmitted diseases at the beginning of the twentieth century. They tested their model with real data of the spread of plague in Bombay between 1905 and 1906. They divided the population into three compartments, S , for susceptible, I for infected the and R for removed, the model contains only two parameters : β the infection rate and γ the rate of recovery, whose values of are determined from the observed data. The size of the total population is assumed to be constant :

$$N = S(t) + I(t) + R(t)$$

Their model can be described by the following set of differential equations

$$\begin{cases} S'(t) = -\beta I(t) S(t) \\ I'(t) = \beta I(t) S(t) - \gamma I(t) \\ R'(t) = \gamma I(t) \end{cases} \quad (1.8)$$

The general hypothesis adopted is that in the time interval dt , a fixed proportion of subjects can no longer be considered as "infected". This applies to the proportion of individuals removed

from the infection process because either they are immune, isolated, or dead. This proportion represented by the parameter γ may be zero and the model reduces to an SI model. It may also be equal to 1, in which case all infected are removed from the infection process during the time interval dt . Generally, the recovery rate γ takes an intermediate value.

1.2.2 Analysis of the model

System 1.8 together with the initial condition $S(0) = S_0$, $I(0) = I_0$, and $R(0) = 0$ such that $I_0 = 0$ and $S_0 = N$ is well-posed. Hereafter we consider a susceptible population in which a small number of infectives is introduced. We show that the positive cone,

$$\mathbb{R}_+^3 = \{(S, I, R) \in \mathbb{R}_+^3 \mid S \geq 0, I \geq 0, R \geq 0\}$$

is positively invariant for system 1.8. As $N = \text{constant} = S(t) + I(t) + R(t)$, the study of system 1.8 is reduced to the study of

$$\begin{aligned} S'(t) &= -\beta I(t) S(t) \\ I'(t) &= \beta I(t) S(t) - \gamma I(t) \end{aligned} \quad (1.9)$$

Consider the following set

$$\Delta = \{(0 \leq S \leq N, 0 \leq I \leq N) \mid 0 \leq S + I \leq N\}$$

Dividing the second equation of 1.9 by $\gamma I(t)$, we have

$$\left\{ \frac{I'(t)}{\gamma I(t)} = \frac{\beta S(t)}{\gamma} - 1 \right. \quad (1.10)$$

Then, for $\frac{\beta S(t)}{\gamma} > 1$ each infective individual infects more than one susceptible, and the disease will spread to an increasing number of individuals. This will reduce the number of susceptibles, $S(t)$ such that $\frac{\beta S(t)}{\gamma} < 1$. The ratio $\frac{\beta}{\gamma}$ can then be interpreted as the number of sufficient contacts to transmit the disease in an infective's entire life of infectiousness.

Multiplying by the fraction of susceptible individuals at each instant, we get the total number of new cases infected by a single infectious individual. Equation 1.10 therefore highlights the importance of initial conditions in the general model, since $\frac{\beta S_0}{\gamma} > 1$, there is likely to be an epidemic, on the contrary, $\frac{\beta S_0}{\gamma} < 1$, only a few individuals will be infected before the epidemic dies out.

Let $\mathcal{R}_0 = \frac{\beta N}{\gamma}$. This threshold $\mathcal{R}_0$ is called the basic reproductive rate, it refers to the number of secondary cases produced by an average infectious individual during the entire period of his infectivity in a population that is entirely susceptible.

In case $\mathcal{R}_0 > 1$, it is possible to solve system 1.8 explicitly, and the following properties also hold

1. S and I have the limits at $+\infty$. In effect, we know that $S'(t) = -\beta I(t) S(t) < 0$ so S is a decreasing function, again $S(t) \geq 0$, or $S(\infty) \geq 0$. Reasoning in a similar manner for R , which is decreasing and bounded, we obtain $R(\infty) \geq 0$ and we deduce the result $I(\infty) = N - S(\infty) - R(\infty)$

2. If $S_0 > 0$ and $I_0 > 0$, then $0 < S(\infty) < S_0$ and $I(\infty) = 0$. Consequently, suppose that $S_0 > 0$, $I_0 > 0$ and $R_0 = 0$ (without loss of generality), otherwise just set $\tilde{N} = N - R_0$. On one hand, for all $t > 0$, $S(t) > 0$, $I(t) > 0$, $R(t) > 0$ and

$$\frac{dS}{dR} = -\frac{\beta}{\gamma} S$$

or

$$S(R) = S_0 e^{-\frac{\beta}{\gamma} R} \geq S_0 e^{-\frac{\beta}{\gamma} N} > 0$$

So $0 < S(\infty) < S_0$.

On the other hand, $\lim_{t \rightarrow \infty} S'(t) = -\beta S(\infty) I(\infty)$ exists. Furthermore, $\lim_{t \rightarrow \infty} S'(t) = 0$. Otherwise if $\lim_{t \rightarrow \infty} S'(t) = \alpha < 0$, then there exists a $T > 0$ such that for all $t > T$, $S'(t) < \frac{\alpha}{2}$, and therefore,

$$S(t) < S_0 + \frac{\alpha}{2} < 0 \quad \text{for } t > -\frac{2S_0}{\alpha}.$$

This would contradict the fact that $S(t) > 0$ for all $t > 0$. So $S(\infty) I(\infty) = 0$ and when $S(\infty) > 0$, then $I(\infty) = 0$

3. The final size of the epidemic is given by $R(\infty) = N - S(\infty)$
 4. We have $\frac{dI}{dS} = -1 + \frac{\gamma}{\beta S}$, and by integration we obtain,

$$I = -S + \frac{\gamma}{\beta} \ln S + C$$

where C is a constant. Or

$$I + S - \frac{\gamma}{\beta} \ln S = C = N - \frac{\gamma}{\beta} \ln N.$$

The maximum size of the epidemic, i.e. I_{max} is given by

$$I_{max} = -\frac{\gamma}{\beta} + \frac{\gamma}{\beta} \ln \frac{\gamma}{\beta} = N - \frac{\gamma}{\beta} \ln N$$

since $S = \frac{\gamma}{\beta}$ in I_{max}

1.3 Transmission of DHF in an Age Structured Population

In this section we consider an age structured model for Dengue, a vector borne disease, like malaria. It is crucial to a model we build later in this thesis.

Pongpunsum et. al. [126] studied an SIR model to describe the transmission of Dengue Hemorrhagic Fever (DHF). They divided the human population into two separate groups, viz; an adult class and juvenile class with a differentiated transmission in the two age classes. The structure in age was motivated by DHF case in Thailand where most cases occurred in children less than 15 years old.

The infection occurs in both classes but infection rate for adults is assumed to be less than that of the children. Since most adults are exposed to DHF infection and develop immunity, many are not aware and they are referred to as the silent population. For the infected children, the antibodies will develop and they become adults with immunity. This means that the adults class composes of the silent population, those who are known to have been infected and those who have never had an infection. All the adults however are represented by S_A .

The transmission from an infectious mosquito is the average of the transmission rates to the different subclasses. The dynamics of the age structured SIR model is given by

$$\left\{ \begin{array}{l} \frac{dS'_J}{dt} = \lambda' - \frac{b\beta_J}{N_T+m} S'_J I'_v - (\mu_h + \delta) S'_J \\ \frac{dR'_J}{dt} = r I'_J - (\mu_h + \delta) R'_J \\ \frac{dI'_J}{dt} = \frac{b\beta_J}{N_T+m} S'_J I'_v - (\mu_h + \delta + r) I'_J \\ \frac{dS'_A}{dt} = \delta (S'_J + I'_J + R'_J) - \epsilon \beta_J \frac{b}{N_T+m} S'_A I'_v - \mu_h S'_J \\ \frac{dR'_A}{dt} = r I'_A - \mu_h R'_A \\ \text{and} \\ \frac{dI'_A}{dt} = \epsilon \beta_J \frac{b}{N_T+m} S'_A I'_v - (\mu_h + r) I'_A \end{array} \right. \quad (1.11)$$

where $S'_{J(A)}$, $I'_{J(A)}$, and $R'_{J(A)}$ are the susceptible juveniles (adults), infected juveniles (adults), and recovered juveniles (adults), respectively; N_T , the total population which is a constant; m , the number of other animals the mosquito can bite; b , the average number of bites a mosquito takes per day; λ , the birth rate; μ_h , the death rate similar for all categories; δ , the rate at which the juveniles pass into adulthood, and r is the rate at which the infected juveniles recover. I'_v is the number of infected mosquitoes; β_J , the probability of the virus surviving in the juvenile after being bitten by an infected mosquito, and $\epsilon\beta_J$ is the probability of the virus surviving in a susceptible adult after an infectious bite. ϵ is the ratio between the probability that an adult becomes infected and the probability that a juvenile becomes infected by an infectious mosquito bite which is assumed to be less than one.

The juvenile population dynamics is given by the equation

$$\frac{dN_J}{dt} = \lambda N_T - (\mu_h + \delta) N_J, \quad (1.12)$$

which was obtained by adding the first three equations of 1.11

The dynamics of the vectors is described by

$$\left\{ \begin{array}{l} \frac{dS'_v}{dt} = A - \frac{b\beta_v}{N_T+m} S'_v (I'_J + I'_A) - \mu_h S'_v \\ \text{and} \\ \frac{dI'_v}{dt} = \frac{b\beta_v}{N_T+m} S'_v (I'_J + I'_A) - \mu_h I'_v \end{array} \right. \quad (1.13)$$

where S'_v and I'_v is the number of susceptible and infected mosquitoes respectively; μ_v , the death rate of the mosquitoes; A , the mosquito carrying capacity of the environment and β_v is the

probability that a dengue virus transmitted to the mosquito from an infected human, from a juvenile or an adult is infectious.

The vector population dynamics is given by

$$\frac{d(S'_v + I'_v)}{dt} = A - \mu_v N_v, \quad (1.14)$$

Pongpunsum et. al. normalized the parameters and eliminated the equations for R_J , R_A , and S_v to study the following set of equations

$$\left\{ \begin{array}{l} \frac{dS_J}{dt} = (\mu_h + \delta)(1 - S_J) - \gamma_h S_J I_v \\ \frac{dI_J}{dt} = \gamma_h S_J I_v - (\mu_h + \delta + r) I_J \\ \frac{dS_A}{dt} = \mu_h - \epsilon \gamma_h S_A I_v - \mu_h S_A \\ \frac{dI_A}{dt} = \epsilon \gamma_h S_A I_v - (\mu_h + r) I_A \\ \frac{dI_v}{dt} = \gamma_{vJ}(1 - I_v)I_J + \gamma_{vA}(1 - I_v)I_A - \mu_v I_v \end{array} \right. \quad (1.15)$$

where

$$\gamma_h = \frac{b \beta_J (A/\mu_v)}{N_T + m}, \quad (1.16)$$

and

$$\gamma_{vJ} = \frac{b \beta_v N_{J(A)}}{N_T + m}. \quad (1.17)$$

By setting the RHS of equation 1.15 to zero, they established that a disease free equilibrium $E_0 = (1, 0, 1, 0, 0)$ and an endemic equilibrium $E_1 = (S_J^*, I_J^*, S_A^*, I_A^*, I_v^*)$ exist. Using linearization method they calculated the value of $\mathcal{R}_0$, and showed, using Routh-Hurwitz criteria, that the disease free equilibrium is locally asymptotically stable when $\mathcal{R}_0 \leq 1$.

The endemic equilibrium analysis was not easy and further simplifications were made. Looking at the data from Bangkok, it was clear that adults have only a small or no chance of becoming sick with the disease. So they set ϵ to zero and, since the adults have developed immunity to the DHF, the two categories I'_A and R'_A would cease to exist. Now in the absence of adults and simplifications as explained above they studied an easier system. This enabled them to study the stability of the endemic equilibrium using linearization.

The endemic equilibrium is shown by simulation to be a stable spiral state. The periods of fluctuations in the number of individuals in each class became much shorter in the absence of any age structure. But the spiraling in, is much more severe in the absence of the age structure and the conclusion was that the age structure appears to calm down the fluctuations.

The model is a one patch model and quite different from malaria which attacks both the children and the adults albeit with differentiated magnitude.

1.4 Differential Susceptibility and Infectivity by Hyman

Hyman et al [76], formulated a differential susceptibility and infectivity model for disease transmission. They divided the susceptibles into n groups based on their susceptibilities, and the infectives were divided into m groups according to their infectivities. Using the standard and bilinear incidence for different diseases, they defined explicit formulas for the reproductive number for each subgroup.

The reproductive number for the entire population is a weighted average of the reproductive numbers for the subgroups. They showed that the infection-free equilibrium is globally stable as the reproductive number is less than one for the models with the bilinear incidence or with the standard incidence but no disease-induced death. They then showed that if the reproductive number is greater than one, there exists a unique endemic equilibrium for these models. Numerical examples are provided to demonstrate that the unique endemic equilibrium is asymptotically stable if it exists.

They considered the spread of a disease in a randomly mixing population that approaches a steady state S^0 if there is no disease infection. They assumed that infected individuals become fully immune or are removed from the susceptible population after they recover from the infection thus studying an SIR model.

The susceptibles who are assumed to have differential susceptibilities are divided into n groups $S_1, S_2, \dots, S_n$. The individuals in each group have homogeneous susceptibilities but the susceptibilities of individuals from different groups are distinct. The susceptibles are distributed into the n susceptible subgroups based on their inherent susceptibility, in such a way that the input flow into S_i is $p_i \mu S^0$ with $\sum_{i=1}^n p_i = 1$. The infectives are divided into m groups, $I_1, I_2, \dots, I_m$, such that upon infection, a susceptible individual in group S_i enters group I_j with the probability q_{ij} and stays in this group until recovery or death, where $\sum_{i=1}^n q_{ij} = 1$ for $j = 1, 2, \dots, m$. From this definition the transmission dynamics of infection are governed by the differential equations;

$$\dot{S}_i = \mu (p_i S^0 - S_i) - \lambda_i S_i, \quad i = 1, \dots, n \quad (1.18)$$

$$\dot{I}_j = \sum_{i=1}^n q_{ij} S_i - (\mu + \nu_j) I_j \quad j = 1, \dots, m \quad (1.19)$$

$$\dot{R} = \sum_{j=1}^m \nu_j I_j - (\mu + \delta) R \quad (1.20)$$

μ is the natural death rate in the absence of infection, ν_j is the recovery rate for infectives in group I_j , and δ is the death rate of the recovered or removed individuals.

The basic reproduction number is defined as the largest eigenvalue of the Jacobian matrix at the disease free equilibrium as follows;

For the bilinear incidence we have;

$$R_0 = \sum_{i=1}^n p_i c_o S^0 \alpha_i \sum_{j=1}^m \frac{q_{ij} \beta_j}{\mu + \nu_j} \quad (1.21)$$

For the standard incidence it is defined as

$$R_0 = \sum_{i=1}^n p_i r \alpha_i \sum_{j=1}^m \frac{q_{ij} \beta_j}{\mu + \nu_j} \quad (1.22)$$

They defined the reproductive number of infection in the susceptible group S_i from all infectives for the standard and bilinear incidence model to be

$$R_{0i} = c(S^0) \beta_i \tau_i,$$

where

$$\tau_i := \sum_{j=1}^m \frac{q_{ij}}{\mu + \nu_j}$$

defines the mean duration of infection from all infectives to susceptibles in group S_i and

$$\beta_i := \alpha_i \frac{1}{\tau_i} \sum_{j=1}^m \frac{q_{ij} \beta_j}{\mu + \nu_j}$$

is the mean transmission probability from all infectives to susceptibles in group S_i .

Then the reproductive number for the entire population can be expressed as the weighted average of the group reproductive numbers such that

$$R_0 = \sum_{i=1}^n p_i R_{0i}.$$

Hyman et al, proved that the disease free equilibrium is globally stable when $R_0 < 1$ and unstable otherwise. When $R_0 > 1$ it is shown that there exists an endemic equilibrium and solutions approach this equilibrium asymptotically. They did not proceed to consider the differential susceptibility and infectivity for spatial population.

1.5 A model for Disease transmission in a Metapopulation by J.Arino

Arino et al [8], formulated a model that incorporates the movement of individuals over a range of spatial scales. They investigated the behavior of the system for a case in which the spatial component consists of a ring of patches and the disease is transmitted between different species in multiple patches. A formula for R_0 is provided and a stability analysis for the disease free equilibrium is undertaken.

They formulated a model that involves s species and n spatial patches and they proceeded as follows

Let N_{ip} be the population number of species $i = 1, 2, \dots, n$ in patch $p = 1, 2, \dots, n$. For species i the rate of travel or migration from patch q to patch p is denoted by $m_{ipq} \geq 0$. These rates form a non negative matrix $\hat{M}_i = [m_{ipq}]$ with $m_{ipp} = 0$. They excluded demography and the rate of change of N_{ip} is given by

$$\dot{N}_{ip} = \sum_{q=1}^n m_{ipp} N_{iq} - \Gamma_{ip} N_{ip} \quad (1.23)$$

where $\Gamma_{ip} = \sum_{q=1}^n m_{iqp}$. Setting $N_i = [N_{i1}, N_{i2}, \dots, N_{in}]^T$, where superscript T denotes transpose, gives the travel equation

$$\dot{N}_i = M_i N_i \quad (1.24)$$

with the mobility matrix $M_i = \hat{M}_i - D_i$. Where D_i is a diagonal matrix with Γ_{ip} is the element (p, p) .

From the sign pattern of M_i and the fact that each column sum is 0, it follows that $(-M)$ is a singular M - matrix ([43] Th. 5 :11). The non-zero entries of M_i specify the arcs of a directed graph for travel connections between patches (as vertices). Arino assumed that from each patch p , to patch q there exists a path in the directed graph so that M_i is irreducible.

For each of the n , patches they formulated a general SEIRS compartmental epidemic model for a disease that confers temporary immunity with $S_{ip}, E_{ip}, I_{ip}, R_{ip}$ denoting the Susceptibles, Latent, Infected (Infectious), and Recovered individuals of species i in patch p at time t , respectively. Therefore

$$N_{ip} = S_{ip} + E_{ip} + I_{ip} + R_{ip}.$$

The new borns from species i in patch p enter the susceptible class with the birth term $d_{ip}N_{ip}$. The transmission term between the species is a standard incidence is $\beta_{ijp} \geq 0$ and defines the average number of adequate contacts per unit time in patch p , between susceptible of species i and infectives of species j . Assuming that the rates of travel are independent of disease status and neglecting disease related deaths, they derived the following system of $4sn$ equations for species $i = 1, 2, \dots, s$ in patch $p = 1, 2, \dots, n$

$$\dot{S}_{ip} = d_{ip}(N_{ip} - S_{ip}) + \nu_{ip} R_{ip} - \sum_{j=1}^s \beta_{ij} S_{ip} \frac{I_{ip}}{N_{ip}} + \sum_{q=1}^n m_{ipq} S_{iq} - \Gamma_{ip} S_{ip} \quad (1.25)$$

$$\dot{E}_{ip} = \sum_{j=1}^s \beta_{ij} S_{ip} \frac{I_{ip}}{N_{ip}} - (d_{ip} + \varepsilon_{ip}) + \sum_{q=1}^n m_{ipq} E_{iq} - \Gamma_{ip} E_{ip} \quad (1.26)$$

$$\dot{I}_{ip} = \varepsilon_{ip} E_{ip} - (d_{ip} + \gamma_{ip}) I_{ip} + \sum_{q=1}^n m_{ipq} I_{iq} - \Gamma_{ip} I_{ip} \quad (1.27)$$

$$\dot{R}_{ip} = \gamma_{ip} I_{ip} - (d_{ip} + \nu_{ip}) R_{ip} + \sum_{q=1}^n m_{ipq} R_{iq} - \gamma_{ip} R_{ip} \quad (1.28)$$

To study the stability of this system they proceeded as follows;

Let

$$F = \begin{bmatrix} 0 & G \\ 0 & 0 \end{bmatrix},$$

where $G = \oplus_{k=1}^n G_k$, and $\oplus$ denotes direct sum.

$$V = \begin{bmatrix} A & 0 \\ -C & B \end{bmatrix},$$

with

$$A = \begin{bmatrix} A_{11} & \dots & A_{1n} \\ \dots & \dots & \dots \\ A_{n1} & \dots & A_{nn} \end{bmatrix},$$

and

$$B = \begin{bmatrix} B_{11} & \dots & B_{1n} \\ \dots & \dots & \dots \\ B_{n1} & \dots & B_{nn} \end{bmatrix},$$

and

$$C = \oplus_{k=1}^n C_k$$

Here G_k, A_{jk}, B_{jk}, C_k are $s \times s$ matrices with A_{jk}, B_{jk}, C_k diagonal. Matrix G_k has (i, j) entry equal to $\beta_{ijk} \frac{N_{ik}^*}{N_{jk}^*}$, the (i, i) entry A_{kk} is equal to $d_{ik} + \varepsilon_{ik} + \sum_{l=1}^n m_{ilk}$, whereas for $j = k$ the (i, i) entry of A_{jk} is $-m_{ijk}$.

Matrix B is the same as A but with ε_{ik} replaced by γ_{ik} . Finally C_k has (i, i) entry equal to ε_{ik} .

From the sign pattern and the fact that they are diagonally dominant by columns, A and B are non-singular M -matrices [18], thus A^{-1} and B^{-1} are non-negative. Using the result in [149], a formula was obtained for the basic reproduction number $\mathcal{R}_0$ and the following stability result.

Theorem 1.1. For the model 1.25 with s species and n patches $\mathcal{R}_0 = \rho(GB^{-1}CA^{-1})$, where ρ denotes the spectral radius. If $\mathcal{R}_0 \leq 1$, then the disease free equilibrium is globally asymptotically stable, and if $\mathcal{R}_0 > 1$, then the disease free equilibrium is unstable.

In this study, metapopulation and its dynamical effect on disease transmission has been done but the study did not consider differential susceptibility and differential infectivity in the patches.

1.6 Ross Model

Ronald Ross [128] modeled the transmission of malaria and identified the basic reproduction number as a threshold for the invasion and persistence of infection. The basic reproduction number, which defines the number of secondary infections resulting from one infectious individual in an entirely susceptible population, determines if the disease will invade the population (if it is greater than 1) or die out (if it is less than 1).

The Ross model divides the entire population as either susceptible or infected (hence infectious). Assuming a constant host population H , he defines I_h as the number of hosts that are infectious, thus there are $H - I_h$ susceptible hosts. Using proportions, he used $\frac{I_h}{H}$ and $\frac{H - I_h}{H}$ to define the fraction of infectious and susceptible hosts respectively. The total vector population is similarly defined with V , the total and vector population, and I_v , the number of infectious vectors, such that $S_v = V - I_v$.

According to Ross, an infection only occurs if a susceptible host is bitten by an infectious vector. The rate at which vectors bite is independent of the number of hosts available since it is clear that each vector needs a certain number of blood meals per unit time. Assuming that the hosts are readily available when the vector needs a blood meal, the biting rate will be proportional to the number of vectors and not the number of hosts.

The man biting rate of vectors is taken to be a , and the proportion of infectious bites to a

host, b_1 . Once infected the host will remain infectious for a period $\frac{1}{\gamma}$ and then recover to join again the susceptible class. The probability that a susceptible vector gets infected when it bites an infected host is defined as b_2 . It is assumed that once a mosquito is infected, it remains so for its entire life, which is assumed to be $\frac{1}{\mu}$ time units.

This leads to the pair of equations

$$\begin{cases} \dot{I}_h = b_1 a \frac{(H-I_h)}{H} I_v - (\gamma) I_h \\ \dot{I}_v = b_2 a (V - I_v) \frac{I_h}{H} - \mu I_v \end{cases} \quad (1.29)$$

He found out that the basic reproductive ratio $\mathcal{R}_0$ is a threshold that determines if the malaria will persist or die out. Precisely if

$$\mathcal{R}_0^2 = \frac{a^2 b_1 b_2 V}{\gamma \mu H} \leq 1,$$

malaria will disappear in the long run
and if

$$\mathcal{R}_0^2 = \frac{a^2 b_1 b_2 V}{\gamma \mu H} > 1,$$

malaria will settle to an endemic equilibrium where both the infectious host and infectious vector populations will be positive.

The square on $\mathcal{R}_0$ comes from the fact that two generations are needed for transmission of malaria to take place, first from an infectious host to a susceptible mosquito and then from an infectious mosquito to a susceptible host. The square on the biting rate a , results from the two cycles of biting necessary for an infection to occur. The expression of basic reproduction number shows that it is directly proportional to the vector population and inversely proportional to the host population. The threshold $\mathcal{R}_0^2 = 1$ determines if the disease will break out or die out.

1.7 Global analysis of a vector-host epidemic model with nonlinear incidence by Li-Ming Cai et. al.

Ming [24] proposed an epidemic model with nonlinear incidence that describe the dynamics of diseases spread by vectors (mosquitoes) with constant human and vector populations. Basing their formulating on the Ross-Macdonald model, they included a Holling Type 2 saturation in the vector populations, which is divided into susceptible and infected vectors.

The dynamics of the human host is described by an SIR model, that is, susceptible, infected and removed compartments. They considered transmission of a vector-borne disease where the transmission is affected by the host behaviour and therefore assumed that the number of infective vectors is proportional to those of the infective human hosts. The result of preventive behaviour in hosts may also limit further transmission of the disease. Another factor that affects transmission is immunity, which they argued enhances transmission. From these facts, they assumed that the infection force of the host population saturates with the infected vectors. The following hypotheses guide the dynamics of their model.

- (H_1) : For host population, the total population under consideration is divided into three classes : $S_H(t)$, $I_H(t)$, and $R_H(t)$, the susceptible, infected and recovered respectively at time t . $N_H(t) = S_H(t) + I_H(t) + R_H(t)$ is the total number of the human population and μK is the recruitment rate of the newborns into the susceptible population. The parameters μ , γ_0 , γ_1 represent the natural death rate, recovery rate and rate of human treatment respectively.
- (H_2) : For the vector population, $S_v(t)$, $I_v(t)$, is the number of susceptible and infective vectors respectively, at time t . The recovery class of mosquitoes is omitted since mosquitoes do not recover from parasite infection. The total vector population thus, is given by $N_v(t) = S_v(t) + I_v(t)$. The rate of recruitment for the vector is the constant rate Λ and the total deaths in mosquitos occur at a rate $mN_v(t)$, where m is the per capita mortality. The differential equation governing the dynamics of the vector population is

$$N'_v(t) = \Lambda - mN_v(t).$$

As $t \rightarrow \infty$ the solution N_v approaches Λ/m .

- (H_3) : Let β_1 be the transmission rate from vector to human and β_2 , the transmission from human to vector. Then considering the saturation in the force of infection, the incidence terms for the host and vector populations are given by

$$\frac{\beta_1 S_H I_v}{1 + \alpha_1 I_v}, \quad \frac{\beta_2 S_v I_H}{1 + \alpha_2 I_v},$$

respectively, where α_1 , α_2 determine the level at which the force of infection saturates.

The dynamics of the vector-host epidemics is described by the following equations

$$\begin{aligned} S'_H &= \mu K - \frac{\beta_1 S_H I_v}{1 + \alpha_1 I_v} - \mu S_H \\ I'_H &= \frac{\beta_1 S_H I_v}{1 + \alpha_1 I_v} - (\mu + \gamma) I_H, \\ R'_H &= \gamma I_H - \mu R_H, \\ S'_v &= \Lambda - \frac{\beta_2 S_v I_H}{1 + \alpha_2 I_v} - m S_v \\ I'_v &= \frac{\beta_2 S_v I_H}{1 + \alpha_2 I_v} - m I_v \end{aligned} \tag{1.30}$$

where $\gamma = \gamma_0 + \gamma_1$. Since $N_v \rightarrow \Lambda/m$, then we can assume $S_v = \Lambda/m - I_v$; The system is feasible in the region R_+^5 , the positive orthant of R^5 . Since R_H and S_v can be determined from the total population, the dynamics of model 1.30 is determined by the three dimension nonlinear system

$$\begin{aligned} S'_H &= \mu K - \frac{\beta_1 S_H I_v}{1 + \alpha_1 I_v} - \mu S_H \\ I'_H &= \frac{\beta_1 S_H I_v}{1 + \alpha_1 I_v} - \omega I_H, \\ I'_v &= \frac{\beta_2 S_v I_H}{1 + \alpha_2 I_v} (\Lambda/m - I_v) - m I_v \end{aligned} \tag{1.31}$$

where $\omega = \mu + \gamma$. For the system 1.31, the region

$$\Omega = \{(S_H; I_H, I_v) \in R_+^3 \mid 0 \leq S_H + I_H \leq K, \quad 0 \leq I_v \leq \Lambda/m\}$$

The basic reproduction number for the system was calculated to be

$$\mathcal{R}_0 = \frac{\beta_1 \beta_2 K \Lambda}{m^2 \omega}$$

where and they proved that in the absence of the disease all solutions tend to the disease free equilibrium E_0 , when $\mathcal{R}_0 \leq 1$. When the $\mathcal{R}_0 > 1$, E_0 becomes unstable and an endemic equilibrium emerges in Ω . Linearising the system at the endemic equilibrium and using the second additive matrix result, they established the local and global stability of system 1.12.

Finally they simulated the model and demonstrated some analytical results and explored the significance of the nonlinear incidence. These results showed that even though the stability results do not depend directly on the parameters α_1 and α_2 , a decrease in these two parameters produces an increase in the number of secondary infections. They confirmed therefore that the parameters α_1, α_2 which vary control the magnitude of the infective individuals.

1.8 Ross-Macdonald Model in a Patchy environment by Auger et. al.

Auger et al [11], modified the Ross-Macdonald model and subdivided the populations into n patches with susceptible/infectious humans and vectors residing in the patches. They assumed that some patches can be vector free, the hosts can migrate between the patches but not the vectors, and that the susceptible and infectious individuals have the same dispersal rate.

Basing their equations on the Ross-Macdonald model, they came up with a set of equations describing their new model. They computed the basic reproduction ratio R_0 and proved that when $R_0 \leq 1$, the disease free equilibrium is globally asymptotically stable, and when $R_0 > 1$, there exists an endemic equilibrium, which is also globally asymptotically stable on the biological domain minus the disease free equilibrium.

The transfer rate from patch i to patch j , for $i \neq j$, is denoted by $m_{ij} \geq 0$. The total host population on patch i is denoted N_i . For $i = 1, 2, \dots, n$, the dynamics is given by

$$\dot{N}_i = \sum_{j=1^n, j \neq i} m_{ij} N_j - N_i \sum_{j=1, j \neq i} m_{ij}.$$

This system can be written as

$$\dot{N} = MN \tag{1.32}$$

They used the following notations

- $I_{h,i}$ is the infectious host population on patch i
- p is the number of patches harboring vectors,
- The variables $I_{v,i}$, and V_i are the infectious vector population and constant vector population on patch i , respectively. If $i > p$ there is no vector on patch i , i.e., $V_i=0$
- a is the man biting rate of vectors
- b_1 is the proportion of infectious bites on hosts that produce a patent infection.

- b_2 is the proportion of infectious bites by susceptible vectors on infectious hosts that produce a patent infection.
- μ is the per capita rate of vector mortality.
- γ is the per capita rate of host recovery from infection.

Auger et al numbered the patches such that only the first p patches, $1 \leq p \leq n$, are infested by the vectors. On the patches i for $i > p$, $V_i = 0$, hence $I_{v,i} = 0$

For patches such that $i \leq p$, i.e., where vectors are present, we have

$$\dot{I}_{h,i} = b_1 a I_{v,i} \frac{N_i - I_{h,i}}{N_i} - \gamma I_{h,i} + \sum_{j=i, j \neq i}^n m_{ij} I_{h,j} - I_{h,i} \left(\sum_{j=i, j \neq i}^n m_{ij} \right) \quad (1.33)$$

$$\dot{I}_{v,i} = b_2 a (V_i - I_{v,i}) \frac{I_{h,i}}{N_i} - \mu I_{v,i} \quad (1.34)$$

For $i > p$, there are no vectors on patch i and the equation for the infectious hosts has only the recovery and migration terms. The equation governing the evolution of $I_{h,i}$ is the following

$$\dot{I}_{h,i} = -\gamma I_{h,i} + \sum_{j=i, j \neq i}^n m_{ij} I_{h,j} - I_{h,i} \left(\sum_{j=i, j \neq i}^n m_{ij} \right) \quad (1.35)$$

They used some notations and conventions to represent the whole system as

$$\dot{N} = M N \quad (1.36)$$

$$\dot{I}_h = \beta_1 \text{diag}(N)^{-1} \text{diag}(N - I_h) I_v - \gamma I_h + M I_h \quad (1.37)$$

$$\dot{I}_h = \beta_1 \text{diag}(N)^{-1} \text{diag}(N - I_h) I_v - \gamma I_h + M I_h \quad (1.38)$$

$$\dot{I}_v = \beta_2 \text{diag}(\pi N)^{-1} \text{diag}(\pi(V - I_v)) \pi I_h - \mu \pi I_v \quad (1.39)$$

$$\pi \dot{I}_v = \beta_2 \text{diag}(\pi N)^{-1} \text{diag}(\pi(V - I_v)) \pi I_h - \mu \pi I_v \quad (1.40)$$

where $\text{diag}(X, p, q)$ denotes, the $p \times q$ matrix diagonal matrix whose diagonal is given by the components of X and the other terms are zero for $X \in \mathcal{R}^p$ or $X \in \mathcal{R}^q$

Using the concept of Metzler matrices and with further simplification of the system they showed that the origin is the disease free equilibrium and computed the basic reproduction number as

$$\mathcal{R}_0^2 = \frac{\beta_1 \beta_2}{\mu} \rho(-\text{diag}(\pi m) Z),$$

where $Z = (D - \gamma I)^{-1}(1 : p, 1 : p)$, i.e., the sub-matrix of the p first rows and p first columns of $D - \gamma I$. It was proved that the disease free equilibrium is globally asymptotically stable.

The stability analysis of the endemic equilibria was done using the theory of irreducible monotonic systems, Lyapunov functions and the La Salle invariance principle. They proved that the endemic equilibrium is locally and also globally asymptotically stable. In this study the researchers concentrated on metapopulation of malaria and the dynamics of transmission when the human host migrates. However they did not consider, differentiated susceptibility and infectivity which are important factors in malaria transmission, that we cannot afford to ignore especially in a patchy environment.

Motivated by the above models, we find it necessary to embark on a study for highland malaria in Western Kenya by extending the models reviewed above.

A VECTOR SIS MALARIA MODEL IN A PATCHY ENVIRONMENT WITH AGE STRUCTURE

2.1 Introduction

Malaria is caused by a parasite of a genus plasmodium, that is transmitted by the female anopheles mosquito as it sucks blood from human hosts for its egg development.

In Kenya, malaria is the leading cause of morbidity and mortality. It is estimated that it accounts for 30% of all outpatient attendance and 19% of all admissions to health facilities. The infection ranges from intense in the lowland to endemic in the highlands causing havoc to the public health system especially during periods of suitable climatic conditions. About 20% of all deaths in children under five are as a result of malaria infection (Ministry of Health-Kenya, 2006 Report). Although the disease is found in most parts of the country, the lake and coastal region plus the highland epidemic-prone districts have risks of over 20% and 19% respectively, with an estimated population of 19 million people (or 49% of the total Kenyan population).

The Western Kenya highlands bears the greatest impact of the disease with some areas suffering as high as 70% infection rates. Entomological studies in different highland ecosystems indicate that transmission is heterogeneous. The terrain characteristics and topography which processes rain, results to big differences in epidemic magnitudes. A study identified three hydrological systems including the V-shaped valleys, the U-shaped valleys and the plateaus.

The types of valleys determine drainage quality, the rainfall threshold and consequently vector breeding. The U-shaped valleys are broad shaped and have slow moving rivers or streams with poor drainage. This favors mosquito population growth resulting in high malaria infection rates and incidence.

The V-shaped valleys have narrow bottoms with fast flowing rivers or streams with good drainage, this makes it hard for mosquito to find suitable breeding sites and thus lower their population growth. The plateaus are flat, though they have good drainage, and their ecosystem behaves like the V-shaped valleys unless there are large water bodies like dams. This implies that the terrain characteristics can modify the level of malaria transmission and the rate of development of immunity as the risk of an epidemic is closely related to the level of immunity of the human population [161].

The V-shaped and the U-shaped valleys are located in different geographical locations, thus

clearly separated, and are apart, so in this case the 'spatialization' is discontinuous. Then the natural tool for modeling in this case is metapopulation.

In Kenya malaria is labeled as a 'travelling disease'. Report from Kibera's Ushirika clinic in Nairobi, where malaria is the main disease treated daily, estimates that 80% percent of the people treated for malaria had travelled out of Nairobi. This implies they got infected and returned, with symptoms only appearing once they are back in the estate. The neighbouring estate is inhabited by people who live near Lake Victoria, where the disease is widespread. By being less exposed to malaria, many lose the semi-immunity they used to have. This is why they contract it easily when travelling upcountry,

(<http://www.irinnews.org/Report/73501/KENYA-Climate-change-and-malaria-in-Nairobi>). This brings the disease to the city which has been free of malaria. Then metapopulation models with migration between the patches suits the 'highland malaria' dynamics in Kenya. Understanding human movement will facilitate the identification of individuals and key zones in malaria transmission and may guide intervention measures, monitoring and improve the intervention programs.

Result from literature also confirms that children between 1 and 5 years represent a very significant source of mosquito infections compared to adults [50] and they are not also bitten in same way as adults [117, 137]. Moreover it is well observed that there is a differentiation in these two age groups as most deaths occur in infants and parasitemia levels of infected individuals decrease with age [146]. This naturally constricts us to differentiated susceptibility and infectivity models.

With the predicted climate change [113], the epidemics may become more frequent and severe posing a serious threat on the already strained public health system. Furthermore, due to the large areas affected by epidemic malaria, it may not be affordable to spray every house with indoor residual insecticides [161]. A model that would lead to a better understanding of transmission and risk of severe disease comes in handy at this time.

Mathematical modeling plays a major role in understanding infectious disease dynamics as it describes the complex disease transmission processes. It also provides insights into the disease dynamics during the rigorous analysis process. The first malaria model was formulated by Sir. Ronald Ross in 1911 [128]. Since then, modeling of vector borne disease has grown in leaps and bounds as shown in reviews [17, 34, 37, 61]. Hyman [76] et al. formulated a general differential susceptibility and differential infectivity model for a disease transmission. They proved that the disease free equilibrium is globally stable when $\mathcal{R}_0 \leq 1$ and unstable otherwise. When $\mathcal{R}_0 > 1$ they show that there exists an endemic equilibrium and solutions approach this equilibrium asymptotically. The model can only be used for directly transmitted disease. Pongsumpun [126] formulated a model to study the influence of age structure in an SIS model for Dengue Hemorrhagic Fever (DHF) and showed that age structure reduces the periods of oscillation on the susceptible and infected human population and the infected mosquito population. It also tightens the spiraling into the endemic equilibrium state. The difference with our intended model is the metapopulation setting, the differentiated patch and age susceptibility and infectivity, not forgetting the disease malaria.

Auger et.al. [11] modified Ross model to n patches. A thorough analysis on the local and global stability of the disease free and the endemic equilibrium is done using $\mathcal{R}_0$. The model however assumed that the susceptibility and infectivity is similar in all the patches, an important difference that we wish to model considering the heterogeneity in the ecosystem of Western Kenyan Highlands.

Motivated by the work of [11] and [126], we formulate an age structured malaria model with susceptibility and infectivity of an individual depending on their age and residence patch. We shall then compare our results with the current findings.

The region is subdivided into homogenous patches and the population in the patches is further subdivided into children, ages 1-5, and adults, over 5 years of age. The susceptibility of the two age groups is differentiated to depend on the patch where the individual resides. The two age groups are allowed to migrate between the patches, making short visits to return to their residences later.

We assume that the rate of migration does not depend on the epidemiological status and age of the human host, that is, that the infected or uninfected children move with the same migration rate as the infected or uninfected adults. We also assume that mosquitoes migrate between some patches, when these patches are not far apart.

In Section 2, we formulate a mathematical model describing the dynamics of an age structure of the human population occupying the patches with migration rates between the patches similar for the two classes. In Section 3, the system is reduced to a more compact form for ease of analysis. In Section 4, we describe the basic properties of the model. In Section 5, we study the stability of the model. An example of the model in two patches is presented in Section 6 and numerical analysis is done in Section 7.

2.2 The model

Let S_{hi}^C (S_{hi}^A), denote the susceptible children (adults) in patch i respectively, while I_{hi}^C (I_{hi}^A) denotes the infective children (adults) respectively. The vector population is likewise identified by S_{vi} and I_{vi} for the susceptible and infective vectors respectively. The total human population which is assumed to be constant, is given by the sum of all host in all the patches as $N_h = S_h + I_h$. So that $N_i = N_i^C + N_i^A$ denotes the total host population in patch i . The total vector population is given by the sum of all vectors in all the patches as $V = S_v + I_v$ and $V_i = S_i + I_i$ is the total vector population in patch i . Migration of hosts is allowed in and out of the patch with an assumption that those who visit return to their resident patches (i.e. they make short visits sufficient for transmission to take place). We assume a uniform migration rate $m_{ij} \geq 0$, $i = 1..n$ denoting proportion of individuals migrating from patch j to patch i . The flow of individuals into the different compartments is represented by Diagram 2.1.

The total host population on patch i is defined as $N_i = N_i^C + N_i^A$ and its dynamics is given by $\dot{N}_i = \dot{N}_i^C + \dot{N}_i^A$, and

$$\begin{aligned}\dot{N}_i^C &= \sum_{j=1, j \neq i}^n m_{ij}^C N_j^C - \sum_{j=1, j \neq i}^n m_{ji}^C N_i^C \\ \dot{N}_i^A &= \sum_{j=1, j \neq i}^n m_{ij}^A N_j^A - \sum_{j=1, j \neq i}^n m_{ji}^A N_i^A\end{aligned}$$

where $\sum_{j=1, j \neq i}^n N_j$ is the total number of residents of patch i at time t [7].

We define the vector of $\mathbb{R}_+^n$, N^C , N^A and V whose components are respectively N_i^C , N_i^A and V_i .

If we define the matrix M^C by

$$M_{i,j}^C = \begin{cases} m_{ij}^C & \text{if } i \neq j \\ m_{ii}^C = - \sum_{j=1, j \neq i}^n m_{ji}^C & \text{otherwise} \end{cases}$$

and the matrix M^A analogously, the preceding equations are

$$\dot{N}^C = M^C N^C \quad \dot{N}^A = M^A N^A$$

We can also assume that there is some migration for the mosquitoes, at least for some nearby patches. Then in the same way, we introduce the total population of mosquitoes V , with the dynamics

$$\dot{V} = M^v V$$

If all the patches are sufficiently distant from each others, then no migration of mosquitoes occurs then $M^v = 0$

Therefore the dynamics of the total host and vector population on patch i is be defined by

$$\dot{N}_i^\Delta = \mathcal{M} N_i^\Delta \tag{2.1}$$

where N^Δ is the vector defined by

$$N^\Delta = \begin{bmatrix} N^C \\ N^A \\ V \end{bmatrix}$$

and the matrix $\mathcal{M}$ is the diagonal block matrix

$$\mathcal{M} = \begin{bmatrix} M^C & 0 & 0 \\ 0 & M^A & 0 \\ 0 & 0 & M^v \end{bmatrix}$$

Our migration equation becomes

$$\dot{N} = \mathcal{M} N. \tag{2.2}$$

The following are the notations used in the model :

- Λ_i is the recruitment of new born children in patch i .
- $S_{h,i}$, $I_{h,i}$ is the susceptible, infectious host population on patch i , respectively.
- $S_{v,i}$, $I_{v,i}$ is the susceptible, infectious vector population on patch i , respectively.
- a_i is the man biting rate of vectors in patch i .
- b_1 is the proportion of infectious bites on a host that produce a patent infection
- b_2 is the proportion of bites by susceptible vectors on an infected host that produce a patent infection
- $\mu_{h,i}^C(\mu_{h,i}^A)$ is the per capita rate of children (adult) mortality in patch i .
- ν is the rate children translate to adults. Then $\frac{1}{\nu}$ is the mean duration of childhood.
- $\mu_{v,i}$ is the per capita rate of vector mortality.

- γ_i is the per capita rate of host recovery from infection, and $\frac{1}{\gamma_i}$ is the mean duration in the infectious compartment in patch i .

The total host and vector population in all the patches is constant. We assume that the parameter a_i is dependent on the host age group in the patches. We also assume that recovery rate γ_i does not depend on age [146] but on how quickly the host receives treatment and their history of malaria infection. This clearly depends on the nature of the health system in the patch i .

We consider age structured system where children translate to adults at a rate ν .

For the patches $i = 1, \dots, n$ we have the $6n$ equations

$$\begin{aligned}
 \dot{S}_{h,i}^C &= \Lambda_i - b_1 a_i^C \frac{S_{h,i}^C}{N_i} I_{v,i} + \gamma_i I_{h,i}^C - (\mu_i^C + \nu) S_{h,i}^C + \sum_{j=1, j \neq i}^n m_{ij}^C S_{h,j}^C - S_{h,i}^C \sum_{j=1, j \neq i}^n m_{ji}^C \\
 \dot{S}_{h,i}^A &= \nu S_{h,i}^C - b_1 a_i^A \frac{S_{h,i}^A}{N_i} I_{v,i} + \gamma_i I_{h,i}^A - \mu_i^A S_{h,i}^A + \sum_{j=1, j \neq i}^n m_{ij}^A S_{h,j}^A - S_{h,i}^A \sum_{j=1, j \neq i}^n m_{ji}^A \\
 \dot{I}_{h,i}^C &= b_1 a_i^C \frac{S_{h,i}^C}{N_i} I_{v,i} - (\gamma_i + \mu_i^C + \nu) I_{h,i}^C + \sum_{j=1, j \neq i}^n m_{ij}^C I_{h,j}^C - I_{h,i}^C \sum_{j=1, j \neq i}^n m_{ji}^C \\
 \dot{I}_{h,i}^A &= b_1 a_i^A \frac{S_{h,i}^A}{N_i} I_{v,i} + \nu I_{h,i}^C - (\gamma_i + \mu_i^A) I_{h,i}^A + \sum_{j=1, j \neq i}^n m_{ij}^A I_{h,j}^A - I_{h,i}^A \sum_{j=1, j \neq i}^n m_{ji}^A \\
 \dot{S}_{v,i} &= \Lambda_v - \frac{S_{v,i}}{N_i} (b_2 a_i^C I_{h,i}^C + b_2 a_i^A I_{h,i}^A) - \mu_{v,i} S_{v,i} + \sum_{j=1, j \neq i}^n m_{ij}^v S_{v,j} - S_{v,i} \sum_{j=1, j \neq i}^n m_{ji}^v \\
 \dot{I}_{v,i} &= \frac{S_{v,i}}{N_i} (b_2 a_i^C I_{h,i}^C + b_2 a_i^A I_{h,i}^A) - \mu_{v,i} I_{v,i} + \sum_{j=1, j \neq i}^n m_{ij}^v I_{v,j} - I_{v,i} \sum_{j=1, j \neq i}^n m_{ji}^v
 \end{aligned} \tag{2.3}$$

The term Λ_i represents the recruitment of newborns into the susceptible children's class, while $b_1 a_i^C I_{v,i} \frac{(S_{h,i}^C)}{N_i}$ in the first equation represents the differentiated infection of susceptible children by infectious mosquitoes using frequency dependent transmission and a non constant host population in patch i . The term $-\gamma_i I_{h,i}^C$ defines the recovery of infected children in patch i . The term μ_i^C is the per capita death rate of children in patch i , ν is the rate children pass into adulthood, and the last term defines host migration. For the second and third equations the terms are defined similarly for Adults. In the sixth equation, for infectious vectors, the first term corresponds to the infection of susceptible mosquitoes by infected children and infected adults, while the last term $-\mu_i I_{v,i}$ caters for vector mortality.

We will assume in the sequel that $\nu < \gamma_i^A$ for any index i

The rationale for this assumption is that the rate of recovering, for an adult or a child, is considerably greater than the mean sojourn time in the compartment of childhood. Actually

$$\nu \approx \frac{1}{5 \times 365} j^{-1} \text{ and } \gamma_i^A \approx \frac{1}{2 \times 30} j^{-1}$$

The complete system is given by equations (2.3).

2.3 Reduced System

It turns out that system given by equations (2.3) can be rewritten in a triangular form so we need the following theorem to reduce such a system and thus study a smaller system.

Theorem 2.1 (Vidyasagar). Consider the following $\mathcal{C}^1$ system :

$$\begin{cases} \dot{x} = f(x) & x \in \mathbb{R}^n, y \in \mathbb{R}^m \\ \dot{y} = g(x, y) \\ \text{with an equilibrium point, } (x^*, y^*) \text{ i.e} \\ f(x^*) = 0 \text{ and } g(x^*, y^*) = 0 \end{cases} \quad (2.4)$$

If x^* is globally asymptotically stable (GAS) in $\mathbb{R}^n$ for the system $\dot{x} = f(x)$, and if y^* is GAS in $\mathbb{R}^m$, for the system $\dot{y} = g(x^*, y)$, then (x^*, y^*) is (locally) asymptotically stable for (2.4). Moreover if all the trajectories of (2.4) are forward bounded, then (x^*, y^*) is a GAS for (2.4).

To apply the above Vidyasagar theorem we would need to prove the stability analysis of first equation, then we would only have to test the stability of the infection equation.

From equation (2.3) if we add the first and third equation together we get

$$\dot{N}_i^C = \Lambda_{h,i} - (\mu_{h,i}^C + \nu) N_i^C + \sum_{j=1, j \neq i}^n m_{ij}^C N_j^C - \sum_{j=1, j \neq i}^n m_{ji}^C N_i^C$$

Which gives

$$\dot{N}^C = \Lambda_h + (-\text{diag}(\mu_h^C + \nu 1) + M^C) N^C.$$

We note that the matrix $[-\text{diag}(\mu_h^C + \nu 1) + M^C] \geq 0$, is a Metzler matrix (Appendix A.17).

For a matrix M the stability modulus, that we will denote $s(M)$, is the largest real part of the elements of the spectrum $\text{Spec}(M)$ of M

$$s(M) = \max_{\lambda \in \text{Spec}(M)} \text{Re}(\lambda).$$

We have now

$$1^T [-\text{diag}(\mu_h^C + \nu 1) + M^C] = -(\mu_h^C + \nu)^T \ll 0.$$

where $1 = (1, 1, \dots, 1)^T$ This proves that the stability modulus of M^C satisfies

$$s(-\text{diag}(\mu_h^C + \nu 1) + M^C) < 0,$$

implying that this Metzler matrix is non singular, which in turn implies that the opposite its inverse is nonnegative [19] .

Therefore the equilibrium of this linear system is given by

$$\bar{N}^C = -[-\text{diag}(\mu_h^C + \nu 1) + M^C]^{-1} \Lambda_h > 0,$$

and is globally asymptotically stable.

A similar result is obtained for the adult population with an equilibrium denoted by $\bar{N}^A$

$$\bar{N}^A = -\nu (-\text{diag}(\mu^A) + M^A)^{-1} \bar{N}^C > 0,$$

and for the mosquito population with an equilibrium denoted by $\bar{V}$.

$$\bar{V} = -(-\text{diag}(\mu_v) + M^v)^{-1} \Lambda_v > 0.$$

Reduction process

We will now give different expressions for the equation of our system. Depending on the case at hand we will use the most convenient form to give the properties of this system and the corresponding proofs.

We can rewrite system (2.3) in $\mathbb{R}_+^{6n}$, with an immediate variable change

$$\left\{ \begin{array}{l} \dot{N}^C = \Lambda + (-\text{diag}(\mu_h^C + \nu 1) + M^C) N^C \\ \dot{N}^A = \nu N^C + (-\text{diag}(\mu_h^A) + M^A) N^A \\ \dot{V} = \Lambda_v + (-\text{diag}(\mu_v) + M^v) V \\ \dot{I}_{h,i}^C = \beta_1^C I_{v,i} \frac{(N_i^C - I_{h,i}^C)}{N_i} - (\gamma_i + \mu_h^C + \nu 1) I_{h,i}^C + \sum_{j=1, j \neq i}^n m_{ij}^C I_{h,j}^C - I_{h,i}^C \sum_{j=1, j \neq i}^n m_{ji}^C \\ \dot{I}_{h,i}^A = \beta_1^A I_{v,i} \frac{(N_i^A - I_{h,i}^A)}{N_i} + \nu I_{h,i}^A - (\gamma_i + \mu_i^A) I_{h,i}^A + \sum_{j=1, j \neq i}^n m_{ij}^A I_{h,j}^A - I_{h,i}^A \sum_{j=1, j \neq i}^n m_{ji}^A \\ \dot{I}_{v,i} = \frac{(V_i - I_{v,i})}{N_i} (\beta_2^C I_{h,i}^C + \beta_2^A I_{h,i}^A) - \mu_v I_{v,i} + \sum_{j=1, j \neq i}^n m_{ij}^v I_{v,j} - I_{v,i} \sum_{j=1, j \neq i}^n m_{ji}^v \end{array} \right. \quad (2.5)$$

This system is clearly triangular if we consider the first variables (N^C, N^A, V) . By application of Theorem 2.1, the stability analysis of (2.5) is now reduced to the stability analysis of the system

$$\left\{ \begin{array}{l} \dot{I}_{h,i}^C = \beta_1^C I_{v,i} \frac{(\bar{N}_i^C - I_{h,i}^C)}{\bar{N}_i} - (\gamma_i + \mu_h^C + \nu 1) I_{h,i}^C + \sum_{j=1, j \neq i}^n m_{ij}^C I_{h,j}^C - I_{h,i}^C \sum_{j=1, j \neq i}^n m_{ji}^C \\ \dot{I}_{h,i}^A = \beta_1^A I_{v,i} \frac{(\bar{N}_i^A - I_{h,i}^A)}{\bar{N}_i} + \nu I_{h,i}^A - (\gamma_i + \mu_i^A) I_{h,i}^A + \sum_{j=1, j \neq i}^n m_{ij}^A I_{h,j}^A - I_{h,i}^A \sum_{j=1, j \neq i}^n m_{ji}^A \\ \dot{I}_{v,i} = \frac{(\bar{V}_i - I_{v,i})}{\bar{N}_i} (\beta_2^C I_{h,i}^C + \beta_2^A I_{h,i}^A) - \mu_v I_{v,i} + \sum_{j=1, j \neq i}^n m_{ij}^v I_{v,j} - I_{v,i} \sum_{j=1, j \neq i}^n m_{ji}^v \end{array} \right. \quad (2.6)$$

We set

$$\begin{aligned} \beta_1^C &= (\beta_1^C, \beta_1^C, \dots, \beta_1^C)^T \\ \beta_1^A &= (\beta_1^A, \beta_1^A, \dots, \beta_1^A)^T \\ \beta_2^C &= (\beta_2^C, \beta_2^C, \dots, \beta_2^C)^T \\ \beta_2^A &= (\beta_2^A, \beta_2^A, \dots, \beta_2^A)^T \\ \bar{N} &= \bar{N}^C + \bar{N}^A \\ I_h^C &= (I_{h,1}^C, I_{h,2}^C, \dots, I_{h,n}^C)^T \\ I_h^A &= (I_{h,1}^A, I_{h,2}^A, \dots, I_{h,n}^A)^T \\ I_v &= (I_{v,1}, I_{v,2}, \dots, I_{v,n})^T \\ \gamma &= (\gamma_1, \gamma_2, \dots, \gamma_n)^T \end{aligned}$$

Then equation (2.6) can be written, in a vectorialized way, as

$$\begin{cases} \dot{I}_h^C = \text{diag}(\beta_1^C) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{N}^C - I_h^C) I_v - \text{diag}(\gamma + \mu_h^C + \nu 1) I_h^C + M^C I_h^C \\ \dot{I}_h^A = \text{diag}(\beta_1^A) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{N}^A - I_h^A) I_v + \nu I_h^C - \text{diag}(\gamma + \mu_h^A) I_h^A + M^A I_h^A \\ \dot{I}_v = [\text{diag}(\beta_2^C I_h^C) + \text{diag}(\beta_2^A I_h^A)] \text{diag}(\bar{N})^{-1} (\bar{V}_i - I_{v,i}) - \text{diag}(\mu_v) I_v + M^v I_v \end{cases} \quad (2.7)$$

For another variable change, we set

$$\begin{aligned} x &= \text{diag}(\bar{N}^C)^{-1} I_h^C \\ y &= \text{diag}(\bar{N}^A)^{-1} I_h^A \\ z &= \text{diag}(\bar{V})^{-1} I_v. \end{aligned}$$

Rewriting system (2.7) in terms of x , y and z , we have

$$\begin{cases} \dot{x} = \text{diag}(\beta_1^C) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) \text{diag}(1-x) z - \text{diag}(\gamma + \mu_h^C + \nu 1) x + \tilde{M}^C x \\ \dot{y} = \text{diag}(\beta_1^A) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) \text{diag}(1-y) z - \text{diag}(\gamma + \mu_h^A) y + \nu x + \tilde{M}^A y \\ \dot{z} = \text{diag}(\bar{N})^{-1} [\text{diag}(\beta_2^C) \text{diag}(\bar{N}^C) \text{diag}(x) + \text{diag}(\beta_2^A) \text{diag}(\bar{N}^A) \text{diag}(y)] (1-z) - \text{diag}(\mu_v) z + \tilde{M}^v z, \end{cases} \quad (2.8)$$

where we define the matrices

$$\begin{aligned} \tilde{M}^C &= \text{diag}(\bar{N}^C)^{-1} M^C \text{diag}(\bar{N}^C), \\ \tilde{M}^A &= \text{diag}(\bar{N}^A)^{-1} M^A \text{diag}(\bar{N}^A), \\ \tilde{M}^v &= \text{diag}(\bar{V})^{-1} M^v \text{diag}(\bar{V}). \end{aligned}$$

Finally, we will make a final "vectorization" of the system

$$\mathcal{A} = \begin{bmatrix} \text{diag}(\beta_1^C) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) & 0 & 0 \\ 0 & \text{diag}(\beta_1^A) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) & 0 \\ 0 & 0 & \text{diag}(\bar{N})^{-1} \end{bmatrix},$$

$$\mathcal{B} = \begin{bmatrix} 0 & 0 & I_n \\ 0 & 0 & I_n \\ \text{diag}(\beta_2^C) \text{diag}(\bar{N}^C) & \text{diag}(\beta_2^A) \text{diag}(\bar{N}^A) & 0 \end{bmatrix}$$

$$\mathcal{C} = \begin{bmatrix} \text{diag}(\gamma + \nu \mathbf{1} + \mu_h^C) & 0 & 0 \\ -\nu I_n & \text{diag}(\gamma + \mu_h^A) & 0 \\ 0 & 0 & \text{diag}(\mu_v) \end{bmatrix}$$

$$\tilde{\mathcal{M}} = \begin{bmatrix} \tilde{M}^C & 0 & 0 \\ 0 & \tilde{M}^A & 0 \\ 0 & 0 & \tilde{M}^v \end{bmatrix}$$

We observe that $\tilde{\mathcal{M}} = \Delta^{-1} \mathcal{M} \Delta$ where Δ is the diagonal matrix

$$\Delta = \begin{bmatrix} \text{diag}(\bar{N}^C) & 0 & 0 \\ 0 & \text{diag}(\bar{N}^A) & 0 \\ 0 & 0 & \text{diag}(\bar{V}) \end{bmatrix}$$

$\mathcal{A}$ and $\mathcal{B}$ are nonnegative matrices and we claim that $(-\mathcal{C} + \tilde{\mathcal{M}})$ is a stable Metzler matrix (Appendix A.17).

Indeed the stability modulus of $-\mathcal{C}$ is negative and

$$s(-\mathcal{C}) = -\min(\gamma + \nu \mathbf{1} + \mu_h^C, \gamma + \mu_h^A, \mu_v) < 0$$

where the minimum is taken over the components of the 3 positive vectors.

We have

$$-\mathcal{C} + \tilde{\mathcal{M}} \leq -s(\mathcal{C}) I_{3n} + \tilde{\mathcal{M}}$$

Since the matrices involved are Metzler matrices, this implies the following inequality for the corresponding stability modulus

$$s(-\mathcal{C} + \tilde{\mathcal{M}}) \leq -s(\mathcal{C}) I_{3n} + s(\tilde{\mathcal{M}}) = -s(\mathcal{C}) < 0$$

The relation $s(\tilde{\mathcal{M}})$ follows from $s(\tilde{\mathcal{M}}) = s(\mathcal{M})$ and from $\mathbf{1}^T \mathcal{M} = 0$ which implies by Perron-Frobenius that $s(\mathcal{M}) = 0$.

Using the preceding matrices and the vector $X = (x, y, z)$ we rewrite equation (2.8) in a compact form as

$$\dot{X} = \mathcal{A} \text{diag}(\mathbf{1} - X) \mathcal{B} X + (-\mathcal{C} + \tilde{\mathcal{M}}) X \quad (2.9)$$

This system evolves on the unit cube of $\mathbb{R}^{3n}$.

2.4 Basic properties of the model

For any index i we shall use the classical notations : $x < y$ if $x \leq y$ and $x_i \leq y_i$ for some i and we write $x_i \ll y_i$ if $x_i < y_i$ for all i .

Proposition 1 (Positively invariant set).

The unit cube

$$\mathfrak{K} = \{(x, y, z) \in \mathbb{R}^{3n} \mid 0 \leq x \leq 1; \quad 0 \leq y \leq 1, \quad 0 \leq z \leq 1\}.$$

is positively invariant for system (2.8).

Proof

To show the invariance of the unit cube $\mathfrak{K}$, under the flow of the system (2.8), it suffices to show that each of the faces of the cube cannot be crossed.

On the patch i we have

$$\dot{x}_i = \beta_1^C \frac{\bar{V}_i}{\bar{N}_i} z_i (1 - x_i) - (\gamma_i + \mu_{h,i}^C + \nu) x_i + \sum_{j=1, j \neq i}^n m_{ij}^C \frac{\bar{N}_j^C}{\bar{N}_i^C} x_j - x_i \sum_{j=1, j \neq i}^n m_{ji}^C$$

If $x_i = 0$

$$\dot{x}_i = \beta_1^C \frac{\bar{V}_i}{\bar{N}_i} z_i + \sum_{j=1, j \neq i}^n m_{ij}^C \frac{\bar{N}_j^C}{\bar{N}_i^C} x_j \geq 0.$$

implying that $x = 0$ cannot be crossed from positive to negative.

If $x_i = 1$ we have

$$\dot{x}_i = -(\gamma_i + \mu_{h,i}^C + \nu) + \sum_{j=1, j \neq i}^n m_{ij}^C \frac{\bar{N}_j^C}{\bar{N}_i^C} x_j - \sum_{j=1, j \neq i}^n m_{ji}^C$$

The relation $x_i = 1$ is equivalent to $I_{h,i}^C = \bar{N}_i^C$. But $\bar{N}^C$ is a demographic equilibrium satisfying

$$(-\text{diag}(\mu_h^C + \nu 1) + M^C) \bar{N}^C = -\Lambda$$

Taking this fact into account gives

$$-\Lambda_i = -(\mu_{h,i}^C + \nu) \bar{N}_i^C + \sum_{j=1, j \neq i}^n m_{ij}^C \bar{N}_j^C - \bar{N}_i^C \sum_{j=1, j \neq i}^n m_{ji}^C,$$

or equivalently

$$-(\bar{N}_i^C)^{-1} \Lambda_i = -(\mu_{h,i}^C + \nu) + \sum_{j=1, j \neq i}^n m_{ij}^C \frac{\bar{N}_j^C}{\bar{N}_i^C} - \sum_{j=1, j \neq i}^n m_{ji}^C,$$

which gives in turn, since $x_j \leq 1$

$$\dot{x}_i \leq -\gamma_i - (\bar{N}_i^C)^{-1} \Lambda_i < 0.$$

The equation for y_i in patch i is

$$\dot{y}_i = \beta_1^A \frac{\bar{V}_i}{\bar{N}_i} z_i (1 - y_i) - (\gamma_i^A + \mu_{h,i}^A) y_i + \sum_{j=1, j \neq i}^n m_{ij} \frac{N_j^A}{N_i^A} y_j - y_i \sum_{j=1, j \neq i}^n m_{ji}$$

If $y_i = 0$ then

$$\dot{y}_i = \beta_1^A \frac{\bar{V}_i}{\bar{N}_i} z_i + \sum_{j=1, j \neq i}^n m_{ij} \frac{N_j^A}{N_i^A} y_j \geq 0.$$

if $y_i = 1$ a similar argument, as for the case $x_i = 1$ gives

$$\dot{y}_i \leq -\gamma_i - \nu (\bar{N}_i^A)^{-1} \bar{N}_i^C < 0.$$

Finally the equation for z_i in patch i is given by

$$\dot{z}_i = \beta_2^C \frac{\bar{N}_i^C}{\bar{N}_i} x_i (1 - z_i) + \beta_2^A \frac{\bar{N}_i^A}{\bar{N}_i} y_i (1 - z_i) - \mu_{v,i} z_i + \sum_{j=1, j \neq i}^n m_{ij}^v \frac{\bar{V}_j}{\bar{V}_i} z_j - z_i \sum_{j=1, j \neq i}^n m_{ji}^v$$

If $z_i = 0$, then for any i ,

$$\dot{z}_i = \beta_2^C \frac{\bar{N}_i^C}{\bar{N}_i} x_i + \beta_2^A \frac{\bar{N}_i^A}{\bar{N}_i} y_i + \sum_{j=1, j \neq i}^n m_{ij}^v \frac{\bar{V}_j}{\bar{V}_i} z_j \geq 0,$$

and if $z_i = 1$ a similar argument, as for the cases $x_1 = 1$ and $y_i = 1$ gives

$$\dot{z}_i \leq -(\bar{V}_i)^{-1} \Lambda_{v,i} \leq 0.$$

The proposition is proved.

Proposition 2.4.1.

If the matrix $M^C + M^A + M^v$ is irreducible, then the system (2.9) is strongly monotone in the interior of the positively invariant set $[0, 1]^{3n}$

Proof

We utilise the theory of monotone dynamical systems introduced by [67, 72, 68], developed further in [67] and applied in [11].

System (2.9) is monotone if its Jacobian is a Metzler matrix on the unit cube. The Jacobian of system (2.9) is given by

$$J(X) = \mathcal{A} \text{diag}(1 - X) \mathcal{B} + (-\mathcal{C} + \tilde{\mathcal{M}}) - \mathcal{A} \text{diag}(\mathcal{B}X)$$

The Jacobian $J(X)$ is clearly a Metzler matrix since $0 \leq X \leq 1$, which implies that the system is cooperative in the unit cube.

Next, we show that the Jacobian $J(X)$ is an irreducible matrix in the set $[0, 1]^{3n}$. This will imply strong monotonicity of the system in the interior of the unit cube. In this set the diagonal terms of $\text{diag}(1 - X)$ are positive, the same property is satisfied for $\mathcal{A}$, then the connectivity of the associated graph of $\mathcal{A} \text{diag}(1 - X) \mathcal{B}$ is reduced to consider the connectivity of $\mathcal{B}$.

It is well known that a matrix is irreducible if its associated graph is strongly connected. Then only the off diagonal terms are concerned. Then it is sufficient to prove that the matrix

$$\mathcal{A} \text{diag}(1 - X) \mathcal{B} + (-\mathcal{C} + \tilde{\mathcal{M}})$$

is irreducible. For the associated graph we distinguish three categories of vertices : the vertices corresponding to the x_i (children vertices), y_i (adults vertices), z_i (mosquitoes vertices). Considering the matrix $\mathcal{B}$, it is clear that a vertex x_i is connected to the vertices of the same patch i , i.e, y_i and z_i . In other words on the three vertices of patch i are contained in a strongly connected component. The vertices from a patch is a subgraph and we can use the contraction of each (x_i, y_i, z_i) in the general digraph to obtain a new digraph wich represents movements between the patches. To prove the strong connectedness, it is sufficient to prove that this contracted digraph is strongly connected.

To the matrix $\tilde{\mathcal{M}}^C$ correspond a digraph between the patches. This graph is equivalent to the associated digraph of $\mathcal{M}^C$. Since M^C , M^A and M^v are Metzler matrices, we can consider the matrix $M^C + M^A + M^v$ and its corresponding multigraph on the graph of patches. This oriented multigraph represents actually the circulation of malaria parasites between the patches. Malaria parasites can be transported either by child, adults or mosquitoes.

Since we have assumed that the matrix, our graph is strongly connected, consequently the Jacobian is irreducible. We note that we are in a context of multiple species (i.e., childs, adults and mosquitoes) in a metapopulation model as conceptualized in [6].

Our assumption simply means that the circulation of parasites is strongly connected.




Figure 2.1 – Transfer Diagram for the different epidemiological groups in different patches

STABILITY ANALYSIS

3.1 Basic Reproduction Number, $\mathcal{R}_0$

We use the classical framework defined in [32],[149].

The application $\mathcal{F}(X) = \mathcal{A} \text{diag}(1 - X) \mathcal{B} X$ represents the rate of appearance of new infections in the compartments in the patches.

The function $\mathcal{V}(X) = (-\mathcal{C} + \tilde{\mathcal{M}}) X$ is the rate of transfer of individuals in compartments.

If $\mathcal{F}(X)$ is set to zero, system becomes $\dot{X} = (-\mathcal{C} + \mathcal{M}) X$, which is a linear system, and we have already seen that $(-\mathcal{C} + \mathcal{M})$ is a stable Metzler matrix (Appendix A.17).

Proposition 2.

The basic reproduction number of system (2.9) is

$$\mathcal{R}_0 = \rho \left(-(-\mathcal{C} + \tilde{\mathcal{M}})^{-1} \mathcal{A} \mathcal{B} \right)$$

Proof

This is straightforward since the Jacobian of $\mathcal{F}$ computed at the DFE $X = 0$ is

$$F = \mathcal{A} \mathcal{B},$$

and the Jacobian of $\mathcal{V}(x)$ computed at the DFE is $\mathfrak{V} = (-\mathcal{C} + \tilde{\mathcal{M}})$.

The next generation matrix is then $K = (-FV^{-1}) - (-\mathcal{C} + \tilde{\mathcal{M}})^{-1} \mathcal{A} \mathcal{B}$

■

We can develop the expression of $\mathcal{R}_0$ a little further :

$$F = \mathcal{A} \mathcal{B} = \begin{bmatrix} 0 & 0 & \text{diag}(\beta_1^C) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) \\ 0 & 0 & \text{diag}(\beta_1^A) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) \\ \text{diag}(\mu_v) \text{diag}(\beta_2^C) \text{diag}(\bar{N}^C) & \text{diag}(\mu_v) \text{diag}(\beta_2^A) \text{diag}(\bar{N}^A) & 0 \end{bmatrix}. \quad (3.1)$$

$$\mathfrak{V} = -\mathcal{C} + \tilde{\mathcal{M}} = \begin{bmatrix} -\text{diag}(\gamma + \nu 1 + \mu_h^C) + \tilde{M}^C & 0 & 0 \\ \nu I_n & -\text{diag}(\gamma + \mu_h^A) + \tilde{M}^A & 0 \\ 0 & 0 & -\text{diag}(\mu_v) + \tilde{M}^v \end{bmatrix}. \quad (3.2)$$

Then we can compute the nonnegative matrix $-\mathfrak{V}^{-1}$, which is a lower triangular matrix

$$-\mathfrak{V}^{-1} = \begin{bmatrix} V_{11}^- & 0 & 0 \\ V_{21}^- & V_{22}^- & 0 \\ 0 & 0 & V_{33}^- \end{bmatrix}$$

with

$$\begin{aligned} V_{11}^- &= -(-\text{diag}(\gamma + \nu 1 + \mu_h^C) + \tilde{M}^C)^{-1} \\ V_{21}^- &= \nu (-\text{diag}(\gamma + \mu_h^A) + \tilde{M}^A)^{-1} (-\text{diag}(\gamma + \nu 1 + \mu_h^C) + \tilde{M}^C)^{-1} \\ V_{22}^- &= -(-\text{diag}(\gamma + \mu_h^A) + \tilde{M}^A)^{-1} \\ V_{33}^- &= -(-\text{diag}(\mu_v) + \tilde{M}^v)^{-1} \end{aligned}$$

The next generation matrix is a block matrix

$$K = \begin{bmatrix} 0 & 0 & K_{13} \\ 0 & 0 & K_{23} \\ K_{31} & K_{32} & 0 \end{bmatrix},$$

with

$$\begin{aligned} K_{13} &= -\text{diag}(\beta_1^C) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) (-\text{diag}(\mu_v) + \tilde{M}^v)^{-1} \\ K_{23} &= -\text{diag}(\beta_1^A) \text{diag}(\bar{N})^{-1} \text{diag}(\bar{V}) (-\text{diag}(\mu_v) + \tilde{M}^v)^{-1} \\ K_{31} &= -\text{diag}(\mu_v) \text{diag}(\beta_2^C) \text{diag}(\bar{N}^C) (-\text{diag}(\gamma + \nu 1 + \mu_h^C) + \tilde{M}^C)^{-1} + \\ &\quad \nu \text{diag}(\mu_v) \text{diag}(\beta_2^A) \text{diag}(\bar{N}^A) (-\text{diag}(\gamma + \mu_h^A) + \tilde{M}^A)^{-1} (-\text{diag}(\gamma + \nu 1 + \mu_h^C) + \tilde{M}^C)^{-1} \\ K_{32} &= -\text{diag}(\mu_v) \text{diag}(\beta_2^A) \text{diag}(\bar{N}^A) (-\text{diag}(\gamma + \mu_h^A) + \tilde{M}^A)^{-1} \end{aligned}$$

The block structure of K implies (see [11]) that

$$\mathcal{R}_0^2 = \rho(K_{31} K_{13} + K_{32} K_{23})$$

When $\mathcal{R}_0 < 1$, the DFE is locally asymptotically stable, and if $\mathcal{R}_0 > 1$ the DFE is unstable, see [149, 32].

3.2 Main Result

In this section we establish a global stability result for the DFE when $\mathcal{R}_0 \leq 1$ and a global stability result when $\mathcal{R}_0 > 1$. We have the following theorem

Theorem 3.1.

We consider the system (2.9) with the matrix $M^C + M^A + M^v$ irreducible.

Then

If $\mathcal{R}_0 \leq 1$, then the system (2.9) is globally asymptotically stable at the origin

If $\mathcal{R}_0 > 1$, then there exists a unique endemic equilibrium X^* , which is globally asymptotically stable on $\mathfrak{K} \in [0, 1]^{3n}$.

Démonstration. We recall system (2.9).

$$\dot{X} = \mathcal{A} \operatorname{diag}(1 - X) \mathcal{B} X + [-\mathcal{C} + \tilde{\mathcal{M}}] X$$

The Jacobian at the origin will be given by

$$J(X) = \mathcal{A} \operatorname{diag}(1 - X) \mathcal{B} - \mathcal{A} \operatorname{diag}(\mathcal{B} X) + [-\mathcal{C} + \tilde{\mathcal{M}}]$$

and

$$J(0) = \mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}]$$

To prove the first part of the proposition above we assume that $\mathcal{R}_0 \leq 1$. Since $[-\mathcal{C} + \tilde{\mathcal{M}}]$ is a Metzler matrix which is invertible and $\mathcal{A} \mathcal{B} \geq 0$, this is a regular splitting of J [152].

We know from [143, 149, 151] that

$$\mathcal{R}_0 \leq 1 \iff s(\mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}]^{-1}) \leq 0.$$

From the preceding section we know that the Jacobian J is an irreducible Metzler matrix. So, by Perron-Frobenius theorem, there exists a positive vector $\mathbf{c} \gg 0$, such that

$$(\mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}])^T \mathbf{c} = s(J(0)) \mathbf{c} \leq 0.$$

To prove the global stability of the DFE we consider the Lyapunov function

$$V(X) = \langle \mathbf{c} \mid X \rangle,$$

where $\langle \mid \rangle$, denotes the inner product. From the definition of $\mathbf{c} \gg 0$, this function is actually positive definite in the nonnegative orthant.

We compute the derivative of V along the trajectories of 2.9 and find that it is equivalent to

$$\begin{aligned} \dot{V}(X) &= \langle \mathbf{c} \mid (\mathcal{A} \operatorname{diag}(1 - X) \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}]) X \rangle \\ &\leq \langle \mathbf{c} \mid (\mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}]) X \rangle \end{aligned} \tag{3.3}$$

$$= \langle (\mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}])^T \mathbf{c} \mid X \rangle$$

$$= \langle s(J(0)) \mathbf{c} \mid X \rangle$$

$$= s(J(0)) \langle \mathbf{c} \mid X \rangle \leq 0, \tag{3.4}$$

Since $s(J(0)) \leq 0$ the derivative is non positive. The DFE is stable.

We will prove the asymptotic stability when $\mathcal{R}_0 \leq 1$.

First we consider the case when $\mathcal{R}_0 < 1$. Since we know that $\mathcal{R}_0 < 1$ implies $s(J) < 0$, $\dot{V}$ is negative definite, since $c \gg 0$. This proves the asymptotic stability of the DFE.

When $\mathcal{R}_0 = 1$, we consider the largest invariant set contained in the set

$$\mathcal{L} = \{X \mid \dot{V}(X) = 0\}$$

For such an X we have

$$\begin{aligned} 0 &= \langle \mathbf{c} \mid \mathcal{A} \text{diag}(1-X) \mathcal{B} X + [-\mathcal{C} + \tilde{\mathcal{M}} X] \\ &= \langle \mathbf{c} \mid (\mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}]) X - \mathcal{A} \text{diag}(X) \mathcal{B} X \rangle, \end{aligned}$$

but since $\dot{V}(X) = 0$, we have by the inequalities (3.3) and (3.4),

$$\langle \mathbf{c} \mid (\mathcal{A} \mathcal{B} + [-\mathcal{C} + \tilde{\mathcal{M}}]) X \rangle = 0$$

Hence

$$\dot{V}(X) = -\langle \mathbf{c} \mid \mathcal{A} \text{diag}(X) \mathcal{B} X \rangle.$$

Since $\mathbf{c} \gg 0$ then $\mathcal{A} \text{diag}(X) \mathcal{B} X = 0$ or equivalently $\text{diag}(X) \mathcal{B} X = 0$

Now we show that the largest invariant set in $\mathcal{L}$ is reduced to the origin.

$$\text{diag}(X) \mathcal{B} X = \begin{bmatrix} \text{diag}(x) z \\ \text{diag}(y) z \\ \text{diag}(z) (\text{diag}(\beta_2^C) \text{diag}(\bar{N}^C) x + \text{diag}(\beta_2^A) \text{diag}(\bar{N}^A) y) \end{bmatrix}$$

For any index i we must have, $x_i z_i = 0$ and $y_i z_i = 0$

Suppose $x_i = 0$ since

$$\dot{x}_i = \beta_1^C \frac{\bar{V}_i}{\bar{N}_i} z_i (1 - x_i) - (\gamma_i + \mu_{h,i}^C + \nu) x_i + \sum_{j=1, j \neq i}^n m_{ij}^C \frac{\bar{N}_j^C}{\bar{N}_i^C} x_j - x_i \sum_{j=1, j \neq i}^n m_{ji}^C$$

we have

$$\dot{x}_i = \beta_1^C \frac{\bar{V}_i}{\bar{N}_i} z_i + \sum_{j=1, j \neq i}^n m_{ij}^C \frac{\bar{N}_j^C}{\bar{N}_i^C} x_j = 0$$

Then $z_i = 0$ and $x_j = 0$ for any patch j , with a "children" arc leaving j and entering i .

Since $z_i = 0$ and $x_i = 0$ and since

$$\dot{z}_i = \beta_2^C \frac{\bar{N}_i^C}{\bar{N}_i} x_i (1 - z_i) + \beta_2^A \frac{\bar{N}_i^A}{\bar{N}_i} y_i (1 - z_i) - \mu_{v,i} z_i + \sum_{j=1, j \neq i}^n m_{ij}^v \frac{\bar{V}_j}{\bar{V}_i} z_j - z_i \sum_{j=1, j \neq i}^n m_{ji}^v$$

We have

$$\dot{z}_i = \beta_2^A \frac{\bar{N}_i^A}{N_i} y_i + \sum_{j=1, j \neq i}^n m_{ij}^v \frac{\bar{V}_j}{V_i} z_j = 0$$

Again $y_i = 0$ and $z_j = 0$ for any patch j with a “mosquito” arc leaving j and entering i .

Now $x_i = y_i = z_i = 0$ implies

$$\dot{y}_i = \sum_{j=1, j \neq i}^n m_{ij}^A \frac{\bar{N}_j^A}{N_i^A} y_j$$

Which implies that $y_j = 0$ for any patch with a “adult” arc leaving j and entering i .

Now, since any patch can be reached by a path composed of “children”, “adult” or “mosquito” arcs, this proves that $x_i = y_i = z_i = 0$ for any index.

The proof for the global asymptotic stability of the DFE is concluded from LaSalle’s Invariance Principle [98]. □

To prove the second part of our theorem, when $\mathcal{R}_0 > 1$, we need the following theorem from Hirsch [72].

Theorem 3.2.

Let F be a C^1 vector field in $\mathbb{R}^n$, whose flow ϕ preserves $\mathbb{R}_+^n$ for $t \geq 0$ and is strongly monotone in $\mathbb{R}_+^n$. Assume that the origin is an equilibrium and that all trajectories in $\mathbb{R}_+^n$ are bounded. Suppose the matrix-valued map $DF : \mathbb{R}_+^n \rightarrow \mathbb{R}_+^n \times \mathbb{R}_+^n$ is strictly anti-monotone, in the sense that,

$$\text{if } x < y, \text{ then } DF(x) > DF(y),$$

then either all the trajectories in $\mathbb{R}_+^n \setminus \{0\}$ tend to the origin, or else there is a unique equilibrium $p \in \text{Int } \mathbb{R}_+^n$ ($p \gg 0$) and all the trajectories in $\mathbb{R}_+^n \setminus \{0\}$ tend to p .

For our case we shall consider the positively invariant set $[0, 1]^{3n}$, which is diffeomorphic to the nonnegative orthant $\mathbb{R}_+^{3n}$. Since the faces of the cube of type $x_i = 1$ are repulsive for the vector field associated to (2.9), all the trajectories are bounded in $[0, 1]^{3n}$.

We recall the Jacobian $J(X)$ of system (2.9)

$$J(X) = -\mathcal{A} \text{diag}(\mathcal{B} X) + \mathcal{A} \text{diag}(1 - X) \mathcal{B} + [-\mathcal{C} + \mathcal{M}],$$

If we take any $X_1 < X_2 \in X$, then $\text{diag}(1 - X_2) < \text{diag}(1 - X_1)$. Since the matrix $\mathcal{B}$ has on each row a positive term, and $\mathcal{A}$ is a diagonal matrix with positive terms, we deduce

$$\mathcal{A} \text{diag}(1 - X_2) \mathcal{B} < \mathcal{A} \text{diag}(1 - X_1) \mathcal{B}$$

Considering the structure of $\mathcal{B}$ we have, if $X_1 < X_2$, the relation $\mathcal{B} X_1 < \mathcal{B} X_2$ and consequently $-\mathcal{A} \text{diag}(\mathcal{B} X_2) < -\mathcal{A} \text{diag}(\mathcal{B} X_1)$.

Finally we have $J(X_2) < J(X_1)$, therefore the anti monotone criteria is met.

We will prove that no trajectory tends to the origin.

We have $\mathcal{R}_0 > 1$ which is equivalent to $s(J(0)) > 1$. Then there exists a positive vector $\mathbf{c} \gg 0$ such that

$$J(0)^T \mathbf{c} = (\mathcal{A} \mathcal{B} + [-\mathcal{C} + \mathcal{M}])^T \mathbf{c} = s(J(0)) \mathbf{c}.$$

We consider the Chetaev function on a neighborhood of the origin

$$V(X) = \langle \mathbf{c} | X \rangle$$

An simple computation gives

$$\dot{V}(X) = \langle \mathbf{c} | [s(J(0)) I_n - \mathcal{A} \text{diag}(X) \mathcal{B}] X \rangle$$

Then in a sufficiently small neighborhood ε of the origin, in $[0, 1]^{3n}$, $\dot{V}(X) > 0$. This proves that for $\varepsilon > 0$ sufficiently small, the hyperplane $\{X | \langle \mathbf{c} | X \rangle = \varepsilon\}$ is a barrier for the vector field associated to (2.9). This proves that no trajectory tends to the origin. Then we conclude, by Hirsch theorem, the existence of an attractive endemic equilibrium X^* in the interior of the cube.

To prove stability, we shall compute $J(X^*) X^*$:

$$J(X^*) X^* = -\mathcal{A} \text{diag}(\mathcal{B} X^*) X^* + \mathcal{A} \text{diag}(1 - X^*) \mathcal{B} X^* + [-\mathcal{C} + \mathcal{M}] X^*,$$

Taking into account that X^* is an equilibrium gives

$$\mathcal{A} \text{diag}(1 - X^*) \mathcal{B} X^* + (-\mathcal{C} + \tilde{\mathcal{M}}) X^* = 0,$$

therefore

$$J(X^*) X^* = -\mathcal{A} \text{diag}(\mathcal{B} X^*) X^* \ll 0$$

since $X^* \gg 0$.

We have proved that there exists a vector $X^* \gg 0$ such that for the Metzler matrix $J(X^*)$ we have $J(X^*) X^* \ll 0$. This implies that $J(X^*)$ is Hurwitz [19, ?].

This conclusion of the global asymptotic stability of the endemic equilibrium is a result of the theorem 3.2.

3.3 An Example in Two Patches

In this section we give a result to the case of two patches. We shall use the structure defined in Subsection 3.1.

$$\mathcal{F} = \begin{bmatrix} \beta_1^C I_{v,1} \frac{(N_1^C - I_{h,1}^C)}{N_1} \\ \beta_1^C I_{v,2} \frac{(N_2^C - I_{h,2}^C)}{N_2} \\ \beta_1^A I_{v,1} \frac{(N_1^A - I_{h,1}^A)}{N_1} \\ \beta_1^A I_{v,2} \frac{(N_2^A - I_{h,2}^A)}{N_2} \\ \frac{(V_1 - I_{v,1})}{N_1} (\beta_2^C I_{h,1}^C + \beta_2^A I_{h,1}^A) \\ \frac{(V_2 - I_{v,2})}{N_2} (\beta_2^C I_{h,2}^C + \beta_2^A I_{h,2}^A) \end{bmatrix}, \quad \mathcal{V} = \begin{bmatrix} -(\gamma_1 + \mu_1^C + \nu) I_{h,1}^C + m_{12} I_{h,2}^C - m_{21} I_{h,1}^C \\ -(\gamma_2 + \mu_2^C + \nu) I_{h,2}^C + m_{21} I_{h,1}^C - m_{12} I_{h,2}^C \\ -(\gamma_1 + \mu_1^A) I_{h,1}^A + m_{12} I_{h,2}^A - m_{21} I_{h,1}^A \\ -(\gamma_2 + \mu_2^A) I_{h,2}^A + m_{21} I_{h,2}^A - m_{12} I_{h,1}^A \\ -\mu I_{v,1} + m_{12}^v I_{v,2} - m_{21}^v I_{v,1} \\ -\mu I_{v,2} + m_{21}^v I_{v,1} - m_{12}^v I_{v,2} \end{bmatrix}.$$

The basic reproduction number is given by $\rho(-FV^{-1})$. From our example and at the DFE, this matrix is defined by FV^{-1} which has the values

$$\begin{bmatrix} 0 & 0 & 0 & 0 & \frac{\beta_1^C N_1^C}{\mu_v N_1} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{\beta_1^C N_2^C}{\mu_v N_2} \\ 0 & 0 & 0 & 0 & \frac{\beta_1^A N_1^A}{\mu_v N_1} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{\beta_1^A N_2^A}{\mu_v N_2} \\ \frac{\beta_2^C V_1(\gamma_2 + \tilde{\mu}_2^C + m_{12})}{N_1(D)} & \frac{\beta_2^C V_1 m_{12}}{N_1(D)} & \frac{\beta_2^A V_1(\gamma_2 + \mu_2^A)}{N_1(D)} & \frac{\beta_2^A V_1 m_{12}}{N_1(D)} & 0 & 0 \\ \frac{\beta_2^C V_2 m_{21}}{N_2(D)} & \frac{\beta_2^C V_2(\gamma_1 + \tilde{\mu}_1^C + m_{21})}{N_2(D)} & \frac{\beta_2^A V_2 m_{21}}{N_2(D)} & \frac{\beta_2^A V_2(\gamma_1 + \mu_1^A + m_{21})}{N_2(D)} & 0 & 0 \end{bmatrix}$$

where $D = (\gamma_1 + m_{21})(\gamma_2 + m_{12}) - m_{12}m_{21}$. To get the basic reproduction number we need to solve $\det[\lambda I - J]$ which is a 6×6 matrix. Rewriting the matrix in the form

$$\Theta = \begin{bmatrix} 0I_4 & B \\ A & 0I_2 \end{bmatrix}.$$

The determinant of Θ is given by

$$\begin{aligned} \det(\Theta - \lambda I_6) &= \begin{vmatrix} -\lambda I_4 & B \\ A & -\lambda I_2 \end{vmatrix} = \lambda^4 \begin{vmatrix} I_4 & -\frac{1}{\lambda}B \\ A & -\lambda I_2 \end{vmatrix}, \\ &= \lambda^4 \left| \begin{bmatrix} I_4 & 0 \\ A & I_2 \end{bmatrix} \begin{bmatrix} I_4 & -\frac{1}{\lambda}B \\ 0 & -\lambda I_2 + \frac{1}{\lambda}AB \end{bmatrix} \right|. \end{aligned}$$

Using the properties of determinants we have

$$\det(\Theta - \lambda I_6) = \lambda^4 \det(-\lambda I_2 + \frac{1}{\lambda}AB) = \lambda^2 \det(AB - \lambda^2 I_2) = 0.$$

We see after some calculation, that

$$\mathcal{R}_0^2 = \frac{a_0 + \sqrt{a_0^2 - 4a_1}}{2}$$

where

$$\begin{aligned} a_0 &= (A_{11}B_{11} + A_{13}B_{31} + A_{22}B_{22} + A_{24}B_{42}) \\ a_1 &= [(A_{11}B_{11} + A_{13}B_{31})(A_{22}B_{22} + A_{24}B_{42})] - [(A_{21}B_{11} + A_{23}B_{31})(A_{12}B_{22} + A_{14}B_{42})] \end{aligned}$$

3.4 Simulation

In this section we obtain baseline values for two sites : the U-shaped valleys and V-shaped valleys. We also describe our reasons for using these values and the references, where available. We estimate parameter values from published studies and country-wide data. For migration rates, we pick realistically feasible values from literature. For the human population in our model, we consider the two patches, Umutete and Iguhu for the U-shaped valleys, and, Marani and Fort

Tenan for the V-shaped valleys. From the study on the different ecosystems, the plateaus and the U-shaped valleys ecosystem have the characteristic, such that the results for the V-shaped valleys apply to the plateau ecosystem. This results from the fact that on the plateaus, the terrain is characterised by raised but flat topography with very little stagnant water as the water drains down the rivers, to support breeding places for mosquitoes. The only notable differences is where there are large water bodies like dams and reservoirs. In these cases high mosquito population is likely to survive and hence increase malaria transmission and infection. Some suitable references for our values are [26, 48, 118, 147, 161].

3.4.1 Parameter values

b_1 : For the proportion of infectious bites on susceptible hosts we shall use the value 0.011 for the U-shaped and 0.08 for the V-shaped valley ecosystem .

b_2 : The proportion of infectious bites on susceptible mosquitoes, is different for the two ecosystems due to acquired or partial immunity so we assume that it is five times higher in the V-shaped valleys (0.24) than the higher in the U-shaped valleys (0.048).

a_h^C : The man biting rate for the children is assumed to be 0.52 in the U-shaped valleys and 0.42 in the V-shaped valleys. We have assumed that the rate is higher due to the high mosquito population in the U-shaped valleys.

a_h^A The biting rate for the adults is less than that of the children and we shall use the value 0.15 for the adults the U-shaped valleys and 0.12 in the V-shaped valleys.

γ : Since the rate of recovery depends on how fast diagnosis is done and the effectiveness of treatment we assume here the same value for the two ecosystems at 0.33 or 30 days period of infectiousness.

ν : The rate children become adults is not easy to determine, but since the two ecosystems have similar economic status, we shall use the value 0.00283 as used in literature.

μ_1^C : The natural death rate for children under five death for the U-shaped valleys is estimated at 7.9% in Kenya [147]. This figure may include disease related deaths but the data is not elaborate. In this case we have assumed that this high rate is for the U-shaped valleys and 5.9% for the V-shaped valleys which may be suffering fewer deaths due to malaria.

$\mu_1^A = \mu_2^A$: The national per capita death rate for adults in the two ecosystem is estimated to be 3.5%.

μ_v : The per capita mosquito death rate is estimated at 0.033 from literature.

Λ_h : This is the recruitment by birth to the susceptible human compartment, in the V-shaped valleys 0.07 is assumed to be less than the rate in the U-shaped valleys 0.13.

Data for N_1 , N_2 , and N_v was not exactly available so we decided to use estimate values with the total population being 10000 people so that we have 5000 people on each patch (3500 children and 1500 adults for each ecosystem). The mosquito population likewise was estimated to be 20000 mosquitoes in the U-shaped valleys and 1500 mosquitoes in the V-shaped valleys.

The summary of the parameter values described above is is given in table 3.1 below.

3.5 Numerical simulation

In this section, we calculate the reproductive number, $\mathcal{R}_0$ for model 2.3 for the parameter values given in table 3.1.

Table 3.1 – Parameter values and ranges for system 2.3

Parameter	U-Shaped Valleys	V-shaped Valleys
a^C	0.52	0.42
a^A	0.15	0.12
b_1^C	0.011	0.08
b_1^A	0.011	0.08
b_2^C	0.048	0.24
b_2^A	0.024	0.018
μ_h^C	0.079	0.059
μ_h^A	0.033	0.033
μ_v	0.033	0.033
ν_h^A	0.000283	0.000283
m_{12}^C	0.08	0.08
m_{21}^C	0.08	0.08
m_{12}^A	0.5	0.5
m_{21}^A	0.5	0.5
γ	0.0035	0.0035
Λ_h	0.04	0.04
Λ_v	0.13	0.07

3.5.1 The U-shaped Valley Sites : Iguhu and Umutete, $\mathcal{R}_0 = 5.36$

We show a numerical simulation of the malaria model 2.3 in the figures below using the parameter values given in Table 3.1

When the age structuring is ignored the dynamics of the host population in the U-shaped valleys is represented by Figure 3.2. The disease in the age structured model fades out faster than the unstructured. The steady states also settle to the endemic equilibrium faster in the age structured model.

If there is no spatialization the values for the U-shaped valleys for both ecosystems has host population variation represented by Figure 3.3. The interaction between the patches raises infection rate, so that the disease persists in the total population, while it fades out fast when the patches are isolated.




Figure 3.1 – A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the U-shaped valleys system. The age structure in the populations is clearly shown. In this case $\mathcal{R}_0 = 5.36$




Figure 3.2 – A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the U-shaped valleys ecosystem. A case where there is no age structure in the population. In this case $\mathcal{R}_0 = 5.36$




Figure 3.3 – A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the U-shaped valleys ecosystem. There is no age structure and the two ecosystems are treated as one U-shaped valley ecosystem




Figure 3.4 – A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the V-shaped valleys ecosystem with age structure. In this case $\mathcal{R}_0 = 0.96 < 1$

3.5.2 The V-shaped Valley Sites : Fort Tenan and Marani, $\mathcal{R}_0 = 1.67$

When the age structuring is ignored the variation of the host population in the V-shaped valleys is represented by Figure 3.5.




Figure 3.5 – A numerical simulation of model 2.3 using parameter values defined in Table 3.1 for the V-shaped valleys ecosystem. There is no age structure in the populations. In this case $\mathcal{R}_0 = 0.96 < 1$

3.6 Discussion and Biological implications

We have seen that the U-shaped valleys support high mosquito density which means that there is more mosquito bites and more malaria is transmitted in the villages. The age structure helps us differentiate between a child's infectivity from an adult's infectivity to a susceptible mosquito. It is clear from literature [15] that children are a significant source of mosquito infection compared to adults. In this case we use the values $b_2^C = 0.48$ and $b_2^A = 0.24$. The biting rate for the two age groups differ [117, 137] as children are bitten more than adults we have $a_i^C = 0.52$ while $a_i^A = 0.42$. The other difference captured in the model is death rates for the children (which may include malaria induced deaths) $\mu_h^C = 0.079$ while the adults die at the rate $\mu_h^A = 0.033$, which is also the national per capital death rate for Kenya [147, 165]. Infection from a mosquito to the two age groups is the same in each patch and we have, $b_1^C = b_1^A = 0.011$ in U-shaped valleys and $b_1^C = b_1^A = 0.008$ for the V-shaped valleys. Using these values in MAPLE plus the ones provided in the table gives the value of the basic reproduction number as $\mathcal{R}_0 = 5.36 < 1$ for the U-shaped valleys and the variation of the populations is given in figure 3.1. When age structure

is ignored the dynamics of the population at equilibrium is shown in figure 3.2. We note that the populations settle to the endemic equilibrium faster than in the age-structured than in the unstructured system as shown in figure 3.2.

For the populations in the V-shaped valleys, the same conditions apply for the two age groups as in the U-shaped valleys and from table 3.1, the the infectious population goes to zero in the longterm as $\mathcal{R}_0 = 0.96 < 1$. We also observe that the stable equilibrium is also achieved faster in the structured than the unstructured system.

Our model suggests that to a suitable model for malaria should be one that captures; age structure; differentiated patch or region susceptibility, which depends on the immunity of the inhabiting population; differentiated infectivity, which also depends on the immunity and age of the host population, and integrate the mosquito population dynamics. Intervention then can then be done with guidance from the model.

A more comprehensive characterization of results would have to include other types of patches, that may not be terrain related but have different epidemiological characteristics from the U-shaped and the V-shaped valleys. Such patches could take care of cities like Nairobi, where human migration has transferred malaria, and central Kenya where the cool highland ecosystem is disturbed by creation of dams for irrigation, rice cultivation, climate change and migration of population to the economically endowed part of the county.

3.7 Conclusion

In this chapter we have done a mathematical analysis of an age structured metapopulation model, by extending the Ross-Macdonald model. We have assumed that the two age groups, children (0-5 years) and adults (over 5 years) migrate to neighboring patches and that their epidemiological characteristics are ecosystem dependent.

The importance of age structure in the dynamics of malaria can not be overemphasized as most malaria induced deaths occur in infants and the average parasitemia levels of infected individuals decreases with age. Age structuring is also crucial since the severity of the disease, its infectivity and susceptibility is age specific, not forgetting that, the acquired immune response also changes with age. Adding age structure allows age specific control strategies on to reduce disease prevalence.

We assumed that vectors migrate especially to nearby patches, and the migration parameters for hosts are constant, similar and independent of the compartment. For the compartments that are far apart, the migration of mosquitoes is negligible and is set to zero, since the mosquitoes are only able to fly a few kilometers away.

An explicit formula for $\mathcal{R}_0$ is obtained, which although complex due to the infinite number of patches, can be used to explore the effects of the parameters of the model. This formula will allow theoretical exploration of the options and efficiency of targeted public health intervention policies. The example in the two ecosystems simplifies the expression for $\mathcal{R}_0$, which we use simulate our model with some realistic data from Western Kenya.

Our model is an improvement over SIS type models for malaria transmission based on the Ross-

Macdonald framework as follows

- **Age structure.** In most malaria models the host population is lumped together as host, in our model we recognize the differentiated infectivity and susceptibility of the children and adults. We also establish that for age structured models, the equilibrium states are achieved faster than the unstructured populations. The novelty is especially visible when age specific control is taken into consideration because with our model we can say what happens when a certain control strategy targets children or adults, how that will impact the model dynamics.
- **Different Patches.** Considering the delicate ecosystem in the western Kenya highlands, our model differs from the classical models in this very important aspect. Malaria epidemics are generally site specific and this micro-environment consideration is well captured in our model. Our model recognizes the different epidemiological characteristics observed in sites that are as close as 4KM apart but with 8 fold difference in malaria prevalence. In this case we can use our model to explore the dynamics of malaria in the patches. For example if there is no malaria in the U-shaped valleys, we can say what happens to the disease in the region.
- **Vectorialized model.** Due to the large size of the system, it maybe difficult to solve and analyze the model fully, if it has to make any biological sense. When we use the compact form of the model, we have employed recent and original tools to study the system. Thus despite the large system, decomposing it into several hierarchical systems and using appropriate change of variables we have proved that the final system is monotone thus applied the to theory non-negative matrices and the theory of monotone dynamical systems by Hirsch to establish the local and global stability of the Endemic equilibrium which would otherwise be a challenge.

This model can be extended to include intervention strategies by the Ministry of health in Kenya, through ITNs and IRS. Since the disease cause death especially in children the model can also include disease related death rate in the human population. An important factor which is under investigation is the impact of climate related factors to the resurgent epidemics. So far, we have formulated an analytical and numerical analysis which is a foundation of more research and also applicable other vector borne disease

DYNAMICS OF VECTOR-HOST FEEDING CONTACT RATE WITH SATURATION IN MALARIA

4.1 Introduction

Malaria is an infectious disease caused by a parasite of the genus, *Plasmodium*. It is transmitted between human hosts by female anopheles mosquitoes as they seek blood meal for their eggs development. When a mosquito bites an infected person, a small amount of blood is taken in which contains microscopic malaria parasites. When the mosquito takes its next blood meal, these parasites mix with the mosquito saliva and are injected into the person being bitten and the transmission process is perpetuated.

Malaria constitutes a big health problem especially within sub-saharan Africa and Asia. World Malaria Report 2011, estimates that it causes between 250-260 million infections and more than a million deaths (mostly among children in Africa), annually. In Kenya, reports show that despite the many control strategies to eliminate malaria, it has re-emerged and increased in incidence. The disease continues to wreck havoc on millions especially from the poor countries [136].

Vector abundance in Western Kenya is driven by temperature variation, ecosystem characteristics and human activities. The population varies depending on the site, the season and the species of the vector. Some sites in Western Kenya has 12.7 fold indoor resting densities during the long rainy season (March-June) and 23.3 fold during the dry season (January-March) [118]. This implies that the vector populations is never constant as assumed in many models. On the other hand host population changes due seasons and economic activities, natural deaths and death due to diseases like malaria and migration to urban centers and other regions for greener pastures. For our model to capture the reality of the epidemics in Western Kenya highlands, we assume that the host and mosquito populations change with time.

Most malaria models assume a constant human biting rates in their models, which means that hosts are freely available whenever a mosquito wants to bite, but in practice, this is more of a simplifying assumption. Research shows that for small host population, this rate is proportional to the host population size, and for large host population, it is constant [53, 74]. The feeding cycle of a mosquito involves, host-seeking, feeding, resting, site-seeking, oviposition and host see-

king resumes [89, 120]. The probability of finding a host and successfully obtaining a blood meal depends on many factors. Among them is human avoidance and defensive behaviour [40, 105].

If the mosquito survives this process, and has a successful blood meal, it rests, finds a larval habitat, oviposits and continues host seeking. Since this process drives malaria transmission, it is necessary to address the specific form of the mosquito-human contact process. Arditi [81], argues that the rates of (successful predation) contact between a predator and a prey is most properly a function of the ratios of their proportions. This would fit into malaria mosquito which inhabit homesteads and other areas where human hosts are available, like farms and urban areas [139]. It is also clear that this contact rate does not increase without bound, as the predator-prey ratio increases, this is because once a mosquito is fed, it rests before ovipositing, to resume host seeking and biting again [114].

When the predator-prey ratio value is low, the contact rate will be limited by the predators ability to find the prey, on the other hand if the ratio is high, the contact rate is limited by the predators satiation (desired predation rate) [74, 81]. For malaria, the contact rate takes a similar course where the mosquito bites will increase as a function of host-vector ratio until the ratio reaches a critical level [95].

Saturation models are also not lacking in literature. A cholera model with saturation in the incidence was proposed by Capasso [25]. They argue that when there is a real threat to infection people become cautious and take preventive measures which controls further infection. Heesterbeek [59] formulated a saturated individual contact rate in relationships such as courting and marriage, where they assume that the population mixes randomly. Zu and Ma [166] analyzed a SEIR epidemic model whose latent period is described by delay and included a saturated incidence rate. Zhang and Ma [168] studied a SEIR model with saturation in contact rates and did a thorough analysis of its global dynamics.

In 2010, Ming and Li [24] formulated vector borne disease model, where they argue that increasing the density of the susceptible hosts with respect to infected ones leads to Holling type II saturation on the force of infection of host. In this model the biting rate of vectors and both populations are assumed to be constant. Further the model neglects disease related deaths a very crucial factor in malaria infection.

A model by [41] on dengue with variable human population is formulated and analyzed for both local and global dynamics. Ngwa et al [119] analyzed the stability of a malaria model, with disease deaths, recovery and variable host and vector populations. However they assumed that the biting rate of vectors is constant hence their infection term is the one described in [128]. Realising then the need to predict the dynamics and transmission of malaria with great precision, we are motivated to engage in this study, as we pay particular attention to saturation in mosquito feeding habits and the varying host and vector populations.

The rest of the paper is subdivided as follows; Section two covers vector-host contact with saturation. In section three, the saturated contact process model is formulated. Section 4 is dedicated to the existence of equilibria, while section 5 studies the stability of the Disease Free and the Endemic equilibrium. Finally in section six we give some results on numerical simulation.

4.2 Vector-Host Contact with saturation

Vector-host contact results from the need for mosquitoes to obtain a blood meal for their eggs development. A given vector's biting rate is limited by both host population density and its own feeding frequency [93].

Therefore the per vector biting rate should increase as a function of the host-vector ratio until the ratio reaches a critical threshold, which we denote Q_v , above which, biting rate saturates and the average vector can feed at its preferred rate b_v (contacts per vector per time). Below this threshold, the relative scarcity of hosts constrains the rate at which a vector can feed on the given type of hosts (it must seek other sources).

We assume that an average host can receive bites at a maximum rate b_h beyond which it successfully defends itself against the vector (including leaving the place altogether) [95]. Then this threshold density ratio is given by

$$Q_v = \frac{b_v}{b_h}.$$

There are two ways of modeling the biting rate which increases for small host population and then approaches a maximum for large populations. The first is using a smooth verhulst-type function of the form

$$f_1\left(\frac{N_h}{N_v}\right) = b_v \frac{\frac{N_h}{N_v}}{\frac{N_h}{N_v} + A}$$

here b_v is the preferred biting rate, and A , the population density ratio at which there is 50% saturation, and it measures how soon the saturation occurs. Dietz [35] and Ming [24] have used an equivalent form of saturation [92].


The other way of modeling the vector biting rate saturation is by using a continuous and piecewise function with a 'switch point', Q_v deleanating the boundary between the two ranges as used by kribis in [94, 92]. This function takes the form

$$f_2\left(\frac{N_h}{N_v}\right) = \begin{cases} = b_v \frac{\frac{N_h}{N_v}}{Q_v}, & \frac{N_h}{N_v} < Q_v \\ = b_v, & \frac{N_h}{N_v} > Q_v \end{cases} \quad (4.1)$$

where b_v is the maximum preferred biting rate and Q_v is the threshold density ratio. There are other Holling-type responses suggested in [59]. The difference between the two saturation models is the saturation sharpness, that is, how quickly the per vector biting rate levels off as the hosts become plentiful, that a vector can feed at its preferred rate, b_v . The saturation is gradual for the function f_1 model and sudden for the model described by f_2 see figure 4.1.

The sharpness is crucial for our malaria model because mosquito populations can change drastically in a short span of time [92]. We also choose the predator-prey type or response since the biting in malaria transmission is vector initiated. The fact that the second model is also easy and captures more dynamics adds to the reason we apply it here in this malaria model.

From the reasons above, we shall assume a saturated contact process as used in [92, 94] with the so-called Holling Type 1 form. Under this assumption, the per-vector contact rate can be

Figure 4.1 – A graphical comparison between f_1 and f_2

described as a function of the host-vector density ratio $z = \frac{N_h}{N_v}$ as

$$f(z) = b_v \min\{z/Q_v, 1\}.$$

When $z > Q_v$ (many host per vector) the rate completely saturates at the maximum desired biting rate $f(z) = b_v$, while for $z < Q_v$ (ie. few hosts per vector) $f(z) = b_v \frac{z}{Q_v}$, and the rate rises linearly with the host-vector ratio. The later is our interest in this study since the saturated contact process $f(z) = b_v$ has been used in the classical Ross model for malaria [128]. Substituting $z = \frac{N_h}{N_v}$ in the function $f(z)$ and multiply by N_v , we obtain the total biting rate which is

$$b_v \min(N_h, Q_v N_v).$$

which can be rewritten in the form

$$\min \{b_h N_h, b_v N_v\}.$$

We note that for the current host density the maximum number of vectors that can effectively bite hosts at one given time is $Q_v N_v$. Therefore the parameter Q_v is an important determinant in our model. It determines which of the two population densities is driving the biting contact rate.

To examine the rate of appearance of new malaria infections from the rate of mosquito feeding contacts, we have to take into account the probability of infection resulting from an effective contact where one party (host or vector) is infected with malaria parasite and the other is not. Let π_h be the probability that such a contact between an infected vector and an uninfected host results in infecting the host, π_v as the proportion of blood meal contacts between infected hosts and uninfected vectors which result in an infected vector. Further if S_h, I_h are the susceptible and infectious hosts respectively and S_v, I_v are the susceptible and infectious vectors respectively, the new infections will be given as defined in [95].

– For the Hosts

$$b_h N_h \frac{S_h}{N_h} \frac{I_v}{N_v} \cdot \pi_h = \pi_h b_h S_h \frac{I_v}{N_v}$$

– For the Vectors

$$b_h N_h \frac{S_v}{N_v} \frac{I_h}{N_h} \cdot \pi_v = \frac{\pi_v b_v}{Q_v} I_h \frac{S_v}{N_v},$$

since $b_h = \frac{b_v}{Q_v}$

In many malaria models, saturation has been assumed to be constant [11, 128, 145]. Here we consider the density dependent biting rate, and the populations ratio plays a vital role in the transmission. If the vectors-host ratio is low, the bites are few and hence the probability of transmission reduces too. As the ratio increases the infection will rise as a function of the ratio until it reaches the threshold and it becomes a constant. We wish to model this change in biting rate as the vector-host ratio changes and study its effect on the basic reproduction ratio.

4.3 The Model Equations

The model we derive here is mathematically equivalent to the classical Ross model [128]. The saturation in contact processes will address how the infection rates depend on this ratio. We assume that mosquito has a variable population growth such that $\Lambda_v > \mu_v$. For the human population, we assume a density dependent mortality rate, such that the total population vary with time and is modified by a logistic equation that include disease induced deaths.

A description of the variables and parameters used in the model follows in tables 5.1 and 4.2 respectively.

Table 4.1 – Variables used in the model related to infection contact process

Variable	Definition
S_h	Susceptible Host Population Density
S_v	Susceptible Vector Population Density
I_h	Infectious Host Population Density
I_v	Infectious Vector Population Density
N_h	Total Host Population Density (constant)
N_v	Total Vector Population Density (variable)

Table 4.2 – Parameters used in the model related to infection contact process

Parameter	Definition
Λ_h, Λ_v	Hosts, Vectors density dependent birth rate
π_h	probability of host infection per contact
π_v	probability of vector infection per contact
γ_h	Hosts rate of recovery
Q_v	vector-host ratio above which per-vector biting saturates
b_h	host irritability biting threshold
b_v	preferred (max.) vector feeding rate
$\nu_h,$	disease dependent death rate

The dynamics of our model will be governed by the following set of equations :

$$\left\{ \begin{array}{l} \dot{S}_h = \Lambda_h - \pi_h b_h S_h \frac{I_v}{N_v} + \gamma_h I_h - \mu_h S_h, \\ \dot{I}_h = \pi_h b_h S_h \frac{I_v}{N_v} - (\mu_h + \gamma_h + \nu_h) I_h, \\ \dot{S}_v = \Lambda_v - \frac{\pi_v b_v}{Q_v} I_h \frac{S_v}{N_v} - \mu_v S_v, \\ \dot{I}_v = \frac{\pi_v b_v}{Q_v} I_h \frac{S_v}{N_v} - \mu_v I_v, \\ \dot{N}_h = \Lambda_h - \mu_h N_h - \nu_h I_h \\ \dot{N}_v = \Lambda_v - \mu_v N_v. \end{array} \right. \quad (4.2)$$

The term Λ_h in the susceptible hosts compartment corresponds to a constant recruitment of susceptible hosts by natural birth. The transmission term $-\pi_h b_h S_h \frac{I_v}{N_v}$ corresponds to frequency dependent infection of susceptible hosts by infectious mosquitoes, on infection they move to the infectious compartment. The infected hosts who recover $\gamma_h I_h$ become susceptible again as malaria has no permanent immunity. The last terms $-\mu_h S_h$, $-\mu_h I_h$ represents per capita deaths of the susceptible, infected hosts respectively. In the susceptible mosquito vectors, Λ_v represent the recruitment of susceptible mosquitoes by birth. The term $\frac{\pi_v b_v}{Q_v} S_h \frac{I_v}{N_v}$ corresponds to the transmission of malaria to an susceptible mosquito by and infected host. Both the susceptible and infectious mosquitoes are subject to natural deaths as defined in the terms $-\mu_v S_v$, $-\mu_v I_v$ respectively. Infective period of mosquitoes ends with their death due to their relatively short life-cycle so we do not have recovery or immune term in the vector equations [14, 60].

All the parameters in the model are non negative and the model equations are well posed. For initial values $(S_h, I_h, S_v, I_v, N_h, N_v)$ in $\mathbb{R}_+^6$, the solutions exist and remains in the region for all $t \geq 0$.

In the absence of disease the host population dynamics is given by $\dot{N}_h = \Lambda_h - \mu_h N_h$. In this kind of demographic structure, the total human and mosquito population size $N_h(t)$ approaches a carrying capacity $\frac{\Lambda_h}{\mu_h}$ for any non zero initial population size.

The mosquito population $N_v(t)$ also approaches a carrying capacity $\frac{\Lambda_v}{\mu_v}$. For ease of studying the system, we let setting $\beta_h = \pi_h b_h$ and $\beta_v = \frac{\pi_v b_v}{Q_v}$, and the equation now takes the form

$$\left\{ \begin{array}{l} \dot{S}_h = \Lambda_h - \beta_h S_h \frac{I_v}{N_v} + \gamma_h I_h - \mu_h S_h, \\ \dot{I}_h = \beta_h S_h \frac{I_v}{N_v} - (\mu_h + \gamma_h + \nu_h) I_h, \\ \dot{S}_v = \Lambda_v - \beta_v I_h \frac{S_v}{N_v} - \mu_v S_v, \\ \dot{I}_v = \beta_v I_h \frac{S_v}{N_v} - \mu_v I_v. \\ \dot{N}_h = \Lambda_h - \mu_h N_h - \nu_h I_h \\ \dot{N}_v = \Lambda_v - \mu_v N_v. \end{array} \right. \quad (4.3)$$

which is defined in feasible region (ie. where the model makes biological sense)

$$\Gamma = \{(S_h, I_h, S_v, I_v, N_h, N_v) \in \mathbb{R}_+^6 : S_h \leq N_h, 0 \leq I_h \leq N_h, S_v \leq N_v, 0 \leq I_v \leq N_v, N_h \geq 0, N_v \geq 0\}$$

where $\mathbb{R}_+^6$ denotes the non-negative cone of $\mathbb{R}^6$ including its lower dimensional faces. It is clear that Γ is positively invariant with respect to (4.3). We denote the boundary and the interior of Γ by $\partial\Gamma$ and $\overset{\circ}{\Gamma}$ respectively.

STABILITY ANALYSIS : SATURATION

5.1 Global stability of the Disease Free Equilibrium

We recall the equations of the model and use the relation $S_h = N_h - I_h$ and $S_v = N_v - I_v$, to study the system

$$\left\{ \begin{array}{l} \dot{I}_h = \beta_h (N_h - I_h) \frac{I_v}{N_v} - (\gamma_h + \mu_h + \nu_h) I_h \\ \dot{I}_v = \beta_v I_h \frac{N_v - I_v}{N_v} - \mu_v I_v \\ \dot{N}_h = \Lambda_h - \mu_h N_h - \nu_h I_h \\ \dot{N}_v = \Lambda_v - \mu_v N_v. \end{array} \right. \quad (5.1)$$

5.1.1 A compact positively invariant set

Using barrier theorems (e.g. [21, 127]) we prove that the following set

$$K = \left\{ (I_h, I_v, N_h, N_v) \mid 0 \leq I_h \leq N_h \leq \frac{\Lambda_h}{\mu_h}, \quad 0 \leq I_v \leq N_v \leq \frac{\Lambda_v}{\mu_v} \right\}$$

is a positively invariant compact set for system (5.1). Moreover K is a global attractor on the nonnegative orthant $\mathbb{R}_+^4$

Since the ODE is Lipschitz, it is sufficient to check that the vector field induced by the system is either tangent or entering K on the boundary K .

Clearly we have the following implications

- $N_v = 0 \Rightarrow \dot{N}_v > 0$ and $N_v \geq \frac{\Lambda_v}{\mu_v} \Rightarrow \dot{N}_v \leq 0$;
- $I_h = 0 \Rightarrow \dot{I}_h \geq 0$;
- since $I_v \leq N_v$ we have $I_v = 0 \Rightarrow \dot{I}_v \geq 0$;
- $N_h = 0 \Rightarrow \dot{N}_h > 0$ and $N_h \geq \frac{\Lambda_h}{\mu_h} \Rightarrow \dot{N}_h \leq 0$;

- when $N_v = I_v$ and $N_v \geq \frac{\Lambda_v}{\mu_v}$ we have $\dot{N}_v - \dot{I}_v = \Lambda_v - 2\mu_v N_v \leq 0$;
- when $N_h = I_h$ and $N_h \geq \frac{\Lambda_h}{\mu_h}$ we have

$$\dot{N}_h - \dot{I}_h = \Lambda_h - (2\mu_h + 2\nu_h + \gamma_h) N_h \leq 0.$$

The preceding relations prove that all trajectories tends to K , which ends the proof of our claim. This also implies that all the trajectories are forward bounded.

We denote the demographic equilibria by $N_h^* = \frac{\Lambda_h}{\mu_h}$ and $N_v^* = \frac{\Lambda_v}{\mu_v}$.

5.1.2 Reduction of the model

We remark that in the last equation only the state N_v is appearing. Hence our system is a triangular system. Using Vidyasagar theorem on K ([157], Theorem A.4 given in annex) we can reduce the stability study to the stability of the equivalent system

$$\begin{cases} \dot{I}_h = \beta_h (N_h - I_h) \frac{I_v}{N_v^*} - (\gamma_h + \mu_h + \nu_h) I_h \\ \dot{I}_v = \beta_v I_h \frac{N_v^* - I_v}{N_v^*} - \mu_v I_v \\ \dot{N}_h = \Lambda_h - \mu_h N_h - \nu_h I_h \end{cases} \quad (5.2)$$

This system is considered now on

$$\Omega = \{(I_h, I_v, N_h) \mid 0 \leq I_h \leq N_h \leq N_h^*, \ 0 \leq I_v \leq N_v^*\}$$

A similar argument, as in the preceding section, shows that Ω is a global attractor on the nonnegative orthant $\mathbb{R}_+^3$ for system (5.2).

5.1.3 Basic reproduction ratio

For system (5.2) there exists a disease free equilibrium (DFE), which is $(0, 0, N_h^*)$ in Ω . According to the technique by van den Driessche-Watmough [148], we define the transmission vector and the vector of other transfers defined on the "infected components"

$$\mathcal{F} = \begin{bmatrix} \beta_h (N_h - I_h) \frac{I_v}{N_v^*} \\ \beta_v I_h \frac{N_v^* - I_v}{N_v^*} \end{bmatrix}, \quad \mathcal{V} = \begin{bmatrix} -(\gamma_h + \mu_h + \nu_h) I_h \\ -\mu_v I_v \end{bmatrix}$$

The Jacobian of these vectors computed at the DFE gives

$$F = \begin{bmatrix} 0 & \beta_h \frac{N_h^*}{N_v^*} \\ \beta_v & 0 \end{bmatrix}, \quad V = \begin{bmatrix} -(\gamma_h + \mu_h + \nu_h) & 0 \\ 0 & -\mu_v \end{bmatrix},$$

hence




Figure 5.1 – Domain of study

$$-FV^{-1} = \begin{bmatrix} 0 & \frac{\beta_h}{\mu_v} \frac{N_h^*}{N_v^*} \\ \frac{\beta_v}{\gamma_h + \mu_h + \nu_h} & 0 \end{bmatrix}$$

The spectral radius of $-FV^{-1}$ gives finally

$$\mathcal{R}_0^2 = \frac{\beta_v \beta_h}{(\gamma_h + \mu_h + \nu_h) \mu_v} \frac{N_h^*}{N_v^*}$$

5.1.4 Equilibria

Let $(\bar{I}_h, \bar{I}_v, \bar{N}_v)$ denote an equilibria. We have the following system of equations

$$\beta_h (\bar{N}_h - \bar{I}_h) \frac{\bar{I}_v}{N_v^*} = (\gamma_h + \mu_h + \nu_h) \bar{I}_h \quad (5.3a)$$

$$\beta_v \bar{I}_h \frac{N_v^* - \bar{I}_v}{N_v^*} = \mu_v \bar{I}_v \quad (5.3b)$$

$$\Lambda_h = \mu_h \bar{N}_h - \nu_h \bar{I}_h \quad (5.3c)$$

The following relations are satisfied :

– From (5.3a)

$$\bar{I}_v = \frac{\beta_v \bar{I}_h}{\frac{\beta_v}{N_v^*} \bar{I}_h + \mu_v} = \frac{\beta_v \bar{I}_h}{\beta_v \bar{I}_h + \mu_v N_v^*} N_v^* ; \quad (5.4)$$

– from (5.3c)

$$\bar{N}_h = \frac{\Lambda_h - \nu_h \bar{I}_h}{\mu_h} = N_h^* - \frac{\nu_h}{\mu_h} \bar{I}_h. \quad (5.5)$$

Replacing these values in (5.3a) gives

$$\beta_h \left[N_h^* - \frac{\nu_h}{\mu_h} \bar{I}_h - \bar{I}_h \right] \frac{\beta_v \bar{I}_h}{\beta_v \bar{I}_h + \mu_v N_v^*} = (\gamma_h + \mu_h + \nu_h) \bar{I}_h.$$

We see that if $\bar{I}_h = 0$, then $\bar{I}_v = 0$ and therefore $\bar{N}_h = N_h^*$. In other words, we obtain the DFE. Then we suppose $\bar{I}_h \neq 0$. Then, from the last expression we obtain

$$\begin{aligned} \beta_v \beta_h N_h^* - (\gamma_h + \mu_h + \nu_h) \mu_v N_v^* &= \beta_v \left[(\gamma_h + \mu_h + \nu_h) + \frac{\nu_h}{\mu_h} \right] \bar{I}_h \\ &= (\gamma_h + \mu_h + \nu_h) \mu_v N_v^* (\mathcal{R}_0 - 1) \end{aligned}$$

Therefore

$$\bar{I}_h = (\gamma_h + \mu_h + \nu_h) \mu_v N_v^* \frac{\mathcal{R}_0 - 1}{\beta_h \left[(\gamma_h + \mu_h + \nu_h) + \beta_v \left(\frac{\nu_h + \mu_h}{\mu_h} \right) \right]} \quad (5.6)$$

If $\mathcal{R}_0 > 1$ we have $\bar{I}_h > 0$ and, from relation (5.4), $\bar{I}_v > 0$. From (5.3a) we obtain

$$\bar{N}_h - \bar{I}_h = \frac{(\gamma_h + \mu_h + \nu_h)}{\beta_h} \bar{I}_h \frac{N_v^*}{\bar{I}_v} > 0$$

Now from relation (5.5) we get $N_h^* > \bar{N}_h$ and finally from (5.4), we get $\bar{I}_v < N_v^*$.

In conclusion, if $\mathcal{R}_0 > 1$, we have an unique endemic equilibrium in the interior of Ω .

5.1.5 Global Stability of the DFE

The DFE is on the boundary of Ω .

Theorem 5.1.

The disease-free equilibrium $P_0 = (0, 0, N_h^*)$ of system (5.2) is globally asymptotically stable in the nonnegative orthant if and only if $\mathcal{R}_0 \leq 1$ and is unstable if $\mathcal{R}_0 > 1$.

Démonstration.

When $\mathcal{R}_0 > 1$ the unstability of the DFE is a consequence of the theorem in [33, 148].

To prove the global stability, we consider a Lasalle-Lyapunov function V ([56, 100, 97]) on the positively invariant compact set. By this definition we mean that on Ω , V is continuous and nonnegative. The function V is not positive definite. We define

$$V(I_h, I_v) = \beta_v I_h + (\gamma_h + \mu_h + \nu_h) I_v.$$

We take advantage on the fact that the ODE (5.2) can be written

$$\begin{bmatrix} I_h \\ I_v \\ N_h \end{bmatrix} = \begin{bmatrix} -(\gamma_h + \mu_h + \nu_h) & \beta_h \frac{N_h - I_h}{N_v^*} & 0 \\ \beta_v \frac{N_v^* - I_v}{N_v^*} & -\mu_v & 0 \\ -\nu_h & 0 & -\mu_h \end{bmatrix} \begin{bmatrix} I_h \\ I_v \\ N_h \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \Lambda_h \end{bmatrix}$$

That we can write, in short, $\dot{X} = A(X) X + b$ If we define $v = (\beta_v, (\gamma_h + \mu_h + \nu_h), 0)^T$, then Then the derivative along the trajectories $\dot{V}$ is given by

$$\dot{V} = v^T A(X) X + v^T b$$

But $v^T b = 0$, then $\dot{V} = v^T A(X)$

$$\begin{aligned} v^T A(X) &= \left[-\beta_v (\gamma_h + \mu_h + \nu_h) + (\gamma_h + \mu_h + \nu_h) \beta_v \left[1 - \frac{I_v}{N_v^*} \right], \quad \beta_v \beta_h \frac{N_h - I_h}{N_v^*} - (\gamma_h + \mu_h + \nu_h) \mu_v, \quad 0 \right] \\ &= \left[-(\gamma_h + \mu_h + \nu_h) \beta_v \frac{I_v}{N_v^*}, \quad \beta_v \beta_h \frac{N_h - I_h}{N_v^*} - (\gamma_h + \mu_h + \nu_h) \mu_v, \quad 0 \right] \\ &\leq \left[-(\gamma_h + \mu_h + \nu_h) \beta_v \frac{I_v}{N_v^*}, \quad (\gamma_h + \mu_h + \nu_h) \mu_v (\mathcal{R}_0 - 1), \quad 0 \right] \end{aligned}$$

For the inequality in the last row we assume $N_h - I_h \leq N_h^*$ on Ω .

Finally, with $\mathcal{R}_0 \leq 1$, since

$$\begin{aligned} \dot{V} &= v^T \begin{bmatrix} I_h \\ I_v \end{bmatrix} \\ \dot{V} &\leq -(\gamma_h + \mu_h + \nu_h) \beta_v \frac{I_v}{N_v^*} I_h + (\gamma_h + \mu_h + \nu_h) \mu_v (\mathcal{R}_0 - 1) I_v \leq 0 \end{aligned}$$

Now the largest invariant set contained in the set E defined by

$$E = \{(I_h, I_v, N_h) \in \Omega | \dot{V}(I_h, I_v) = 0\}$$

is certainly contained in the set of points for which $I_v = 0$ or $I_h = 0$. We have two situations :

- To be in an invariant set with $I_v = 0$ has for consequences that $I_h = 0$ and consequently $N_h = N_h^*$, hence to be at the DFE;
- To be in an invariant set with $I_h = 0$ implies either $I_v = 0$, thus to be at the DFE, or $N_h = I_h = 0$, which is cannot be contained in an invariant set by the last equation system (5.2).

We have proved that the largest invariant set contained in $\dot{V} = 0$ and in the positively invariant set Ω is the singleton constituted by the DFE. By Lasalle 's theorem [100, 97] this proves that the DFE is globally asymptotically stable in Ω , hence in the nonnegative orthant. $\square$

5.1.6 Stability of the endemic equilibrium

In this section we will prove that, when $\mathcal{R}_0 > 1$ then the unique endemic equilibrium which is asymptotically stable.

Theorem 5.2.

The endemic equilibrium $EE = (\bar{I}_h, \bar{I}_v, \bar{N}_h)$ of system (5.2), is given by the equations (5.6), (5.4) and (5.5), and is asymptotically stable in Ω .

Démonstration. We consider the Jacobian of system (5.2)

$$J(I_h, I_v, N_v) = \begin{bmatrix} -\beta_h \frac{I_v}{N_v^*} - (\gamma_h + \mu_h + \nu_h) & \beta_h \frac{(N_h - I_h)}{N_v^*} & \beta_h \frac{I_v}{N_v^*} \\ \beta_v \frac{(N_v^* - I_v)}{N_v^*} & -\beta_v \frac{I_h}{N_v^*} - \mu_v & 0 \\ -\nu_h & 0 & -\mu_h \end{bmatrix}$$

We can see that our system is not monotone and cannot be monotone for any orthant of $\mathbb{R}^n$, since we have the (3,1) entry of J negative and the (1,3) entry positive.

The Jacobian computed at the EE, using the relation of (5.3), can be expressed as

$$J(\bar{I}_h, \bar{I}_v, \bar{N}_v) = \begin{bmatrix} -\beta_h \frac{\bar{N}_h}{N_v^*} \frac{\bar{I}_v}{\bar{I}_h} & (\gamma_h + \mu_h + \nu_h) \frac{\bar{I}_h}{\bar{I}_v} & \beta_h \frac{\bar{I}_v}{N_v^*} \\ \mu_v \frac{\bar{I}_v}{\bar{I}_h} & -\beta_v \frac{\bar{I}_h}{\bar{I}_v} & 0 \\ -\nu_h & 0 & -\mu_h \end{bmatrix}$$

We will examine the characteristic polynomial of $J(\bar{I}_h, \bar{I}_v, \bar{N}_v)$, denoted by $\text{Char}(J)(X) = X^3 + a_1 X^2 + a_2 X + a_3$.

It is well known that

$$a_1 = -\text{Trace}(A) \quad a_3 = -\text{Det}(A)$$

The coefficient a_2 is equal to the sum of the principal minors 2×2 .

The Routh-Hurwitz criteria ensures that $J(\bar{I}_h, \bar{I}_v, \bar{N}_v)$ is Hurwitz iff $a_i > 0$ for $i = 1, \dots, 3$ and $a_1 a_2 - a_3 > 0$.

The upper 2×2 - block of J is

$$M = \begin{bmatrix} -\beta_h \frac{\bar{N}_h}{N_v^*} \frac{\bar{I}_v}{\bar{I}_h} & (\gamma_h + \mu_h + \nu_h) \frac{\bar{I}_h}{\bar{I}_v} \\ \mu_v \frac{\bar{I}_v}{\bar{I}_h} & -\beta_v \frac{\bar{I}_h}{\bar{I}_v} \end{bmatrix}$$

Hence the determinant is given by

$$\begin{aligned} \det(M) &= \beta_h \beta_v \frac{\bar{N}_h}{N_v^*} \frac{\bar{I}_h}{\bar{I}_v} \frac{\bar{I}_v}{\bar{I}_h} - \mu_v (\gamma_h + \mu_h + \nu_h) \frac{\bar{I}_h}{\bar{I}_v} \frac{\bar{I}_v}{\bar{I}_h} \\ &= \beta_h \beta_v \frac{\bar{N}_h}{N_v^*} - \mu_v (\gamma_h + \mu_h + \nu_h) \end{aligned}$$

The determinant of J is

$$\begin{aligned}
J(\bar{I}_h, \bar{I}_v, \bar{N}_v) &= -\mu_h \left(\beta_h \beta_v \frac{\bar{N}_h}{N_v^*} - \mu_v (\gamma_h + \mu_h + \nu_h) \right) - \nu_h \beta_h \beta_v \frac{\bar{I}_h}{N_v^*} \\
&= \mu_h \mu_v (\gamma_h + \mu_h + \nu_h) - \beta_h \beta_v \frac{\mu_h \bar{N}_h + \nu_h \bar{I}_h}{N_v^*} \\
&= \mu_h \mu_v (\gamma_h + \mu_h + \nu_h) - \beta_h \beta_v \frac{\Lambda_h}{N_v^*} \\
&= \mu_h \mu_v (\gamma_h + \mu_h + \nu_h) - \beta_h \beta_v \frac{\mu_h N_h^*}{N_v^*} \\
-a_3 &= -\mu_h \mu_v (\gamma_h + \mu_h + \nu_h) (\mathcal{R}_0 - 1) < 0
\end{aligned}$$

We consider now the trace of J which is negative

$$\text{Trace}(J) = -a_1 = -\beta_h \frac{\bar{N}_h}{N_v^*} \frac{\bar{I}_v}{\bar{I}_h} - \beta_v \frac{\bar{I}_h}{\bar{I}_v} - \mu_h < 0$$

We have now to compute a_2 the sum of the 2×2 principal minors .

$$\begin{aligned}
a_2 &= \frac{\bar{N}_h \beta_h \beta_v - \bar{N}_v^* \mu_v (\gamma_h + \mu_h + \nu_h)}{\bar{N}_v^*} + \frac{\bar{I}_v \beta_h (\bar{N}_h \mu_h + \bar{I}_h \nu_h)}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} \\
&= \frac{\bar{N}_h \beta_h \beta_v}{\bar{N}_v^*} - \mu_v (\gamma_h + \mu_h + \nu_h) + \frac{\bar{I}_v \beta_h (N_h^* \mu_h)}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} \\
&= \frac{\bar{N}_h \beta_h \beta_v}{\bar{N}_v^*} - \mu_v \beta_h \frac{(\bar{N}_h - \bar{I}_h) \bar{I}_v}{N_v^* \bar{I}_h} + \frac{\bar{I}_v \beta_h N_h^* \mu_h}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} \\
&= \frac{\bar{N}_h \beta_h \beta_v}{\bar{N}_v^*} - \mu_v \beta_h \frac{\bar{N}_h}{N_v^*} \frac{\bar{I}_v}{\bar{I}_h} + \mu_v \beta_h \frac{\bar{I}_v}{N_v^*} + \frac{\bar{I}_v \beta_h N_h^* \mu_h}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} \\
&= \frac{\bar{N}_h \beta_h \beta_v}{\bar{N}_v^*} - \beta_h \frac{\bar{N}_h}{N_v^*} \frac{1}{\bar{I}_h} \left(\beta_v \bar{I}_h - \beta_v \frac{\bar{I}_h \bar{I}_v}{N_v^*} \right) + \mu_v \beta_h \frac{\bar{I}_v}{N_v^*} + \frac{\bar{I}_v \beta_h N_h^* \mu_h}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} \\
&= \beta_h \beta_v \frac{\bar{I}_h \bar{I}_v}{N_v^*} + \mu_v \beta_h \frac{\bar{I}_v}{N_v^*} + \frac{\bar{I}_v \beta_h N_h^* \mu_h}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} > 0
\end{aligned}$$

The first three requirements of Routh-Hurwitz 's criteria are satisfied. We have now to conduct a long calculation to prove $a_1 a_2 - a_3 > 0$

Let compute $a_1 a_2 - a_3$

$$\begin{aligned}
a_1 a_2 - a_3 &= \left[\beta_h \frac{\bar{N}_h}{N_v^*} \frac{\bar{I}_v}{\bar{I}_h} + \beta_v \frac{\bar{I}_h}{\bar{I}_v} + \mu_h \right] \times \left[\beta_h \beta_v \frac{\bar{I}_h \bar{I}_v}{N_v^*} + \mu_v \beta_h \frac{\bar{I}_v}{N_v^*} + \frac{\bar{I}_v \beta_h N_h^* \mu_h}{\bar{I}_h \bar{N}_v^*} + \frac{\bar{I}_h \mu_h \beta_v}{\bar{I}_v} \right] + \\
&\quad \left[\mu_h \mu_v (\gamma_h + \mu_h + \nu_h) - \beta_h \beta_v \frac{\mu_h N_h^*}{N_v^*} \right] \\
&= \beta_h^2 \beta_v \frac{\bar{N}_h \bar{I}_v^2}{N_v^{*2}} + \beta_h \beta_v^2 \frac{\bar{I}_h^2}{N_v^*} + \mu_h \beta_h \beta_v \frac{\bar{I}_h \bar{I}_v}{N_v^*} + \mu_v \beta_h^2 \frac{\bar{N}_h \bar{I}_v^2}{N_v^{*2} \bar{I}_h} + \mu_h \beta_h \beta_v \frac{\bar{I}_h}{N_v^*} + \\
&\quad \mu_v \mu_h \beta_h \frac{\bar{I}_v}{N_v^*} + \mu_h \beta_h^2 \frac{\bar{N}_h N_h^* \bar{I}_v^2}{N_v^{*2} \bar{I}_h^2} + \mu_h \beta_h \beta_v \frac{N_h^*}{N_v^*} + \mu_h^2 \beta_h \frac{N_h^* \bar{I}_v}{N_v^* \bar{I}_h} + \mu_h \beta_h \beta_v \frac{\bar{I}_h \bar{N}_h}{N_v^*} + \\
&\quad \mu_h \beta_v^2 \frac{\bar{I}_h^2}{\bar{I}_v^2} + \mu_h^2 \beta_v \frac{\bar{I}_h}{\bar{I}_v} + \mu_h \mu_v (\gamma_h + \mu_h + \nu_h) - \mu_h \beta_h \beta_v \frac{N_h^*}{N_v^*} \\
&= \beta_h^2 \beta_v \frac{\bar{N}_h \bar{I}_v^2}{N_v^{*2}} + \beta_h \beta_v^2 \frac{\bar{I}_h^2}{N_v^*} + \mu_h \beta_h \beta_v \frac{\bar{I}_h \bar{I}_v}{N_v^*} + \mu_v \beta_h^2 \frac{\bar{N}_h \bar{I}_v^2}{N_v^{*2} \bar{I}_h} + \mu_h \beta_h \beta_v \frac{\bar{I}_h}{N_v^*} + \\
&\quad \mu_v \mu_h \beta_h \frac{\bar{I}_v}{N_v^*} + \mu_h \beta_h^2 \frac{\bar{N}_h N_h^* \bar{I}_v^2}{N_v^{*2} \bar{I}_h^2} + \mu_h^2 \beta_h \frac{N_h^* \bar{I}_v}{N_v^* \bar{I}_h} + \mu_h \beta_h \beta_v \frac{\bar{I}_h \bar{N}_h}{N_v^*} + \\
&\quad \mu_h \beta_v^2 \frac{\bar{I}_h^2}{\bar{I}_v^2} + \mu_h^2 \beta_v \frac{\bar{I}_h}{\bar{I}_v} + \mu_h \mu_v (\gamma_h + \mu_h + \nu_h) > 0
\end{aligned}$$

This proves the asymptotic stability of the endemic equilibrium. $\square$

5.1.7 Global stability of the endemic equilibrium, when there is no disease induced mortality

We suppose in this section that $\nu_h = 0$. In this case we can reduce the system to a two dimensional system, thanks to Vidyasagar's theorem. The original system is then, for the stability point of view, equivalent to

$$\begin{cases} \dot{I}_h = \beta_h (N_h^* - I_h) \frac{I_v}{N_v^*} - (\gamma_h + \mu_h) I_h \\ \dot{I}_v = \beta_v I_h \frac{N_v^* - I_v}{N_v^*} - \mu_v I_v \end{cases} \quad (5.7)$$

In this case this system is a Ross system, and the global stability of the endemic equilibrium is well known, see for example [12, 100, 167]

5.1.8 Simulation

Using the values in Table 5.1 we simulate system 4.2, and show the dynamics of various populations in the figures below.

5.1.9 Estimating Q_v

The saturation threshold ratio Q_v is an important parameter that determines which of the two populations, that is the host and the mosquito, is driving malaria infection. We will assume that the indoor resting density for mosquitoes is 2.5 for each household that has approximately 6.5 persons (Githeko, personal communication), then $\frac{N_h}{N_v} = 2.6$. We also assume that $b_v = 9$, and $b_h = 10$, this gives $Q_v = 0.9$. Recalling our function for the feeding rate with saturation, $f(z) = b_v \min(\frac{z}{Q_v}, 1)$, implies that $Q_v < z$, and the vector feeds at its preferred rate, b_v . When

Table 5.1 – Parameters used in the model related to infection contact process

Parameter	Value
Λ_h	0.04
Λ_v	0.13
π_h	0.22
π_v	0.48
γ_h	0.33
b_h	3(10)
b_v	9
ν_h	0.0329
μ_h	0.033
μ_v	0.033

$Q_v < z$, N_h must be greater than N_v , so the biting rate will depend on the vector and not the host population. This would result to high vector host contact and high malaria transmission. In this case $\tilde{\mathcal{R}}_0 = 2.495 > 1$ and malaria will persist in the population as shown in figure 5.5. If however the host irritation from vector bites is high, then we assume a $b_h = 3$, then $Q_v = 3 > z$. This implies from our model that there are few host for each vector and the relative scarcity of hosts constrains the rate at which an average vector can feed. The biting rate therefore will be driven by the host and not the vector population. For its survival it may be forced to seek other sources for its blood meal. Reduced host biting translates to lower malaria infection and a reduced basic reproduction number. Then $\tilde{\mathcal{R}}_0 = 0.718 < 1$, the infection dies out, and the infectious population goes to zero as shown in figure 5.4.

5.2 Discussion and Conclusion

The principal motivation in this model has been to capture the non-linear contact process with saturation which is the key factor in the spread of malaria. The natural fluctuations of mosquito populations results in varying host-vector density ratio. We have shown that the two population densities drive the entire infection process through the threshold population density ratio Q_v , which plays a vital role in the basic reproduction number.

The deviation from the constant biting rate addresses the important variability in the vector population especially, which is affected by anti-malaria campaign (like Inside Residual spraying IRS), ecosystem heterogeneity and climatic factors in Western Kenya highlands.

Of importance also is the inclusion of disease induced deaths, having in mind that majority of malaria deaths occur in children in Kenya. Mathematical analysis was done to establish that in the absence of the disease, a disease free equilibrium will always exist if $\tilde{\mathcal{R}}_0 \leq 1$. In the presence of the disease, that is when $\tilde{\mathcal{R}}_0 > 1$, an endemic equilibrium is established with the infectious populations greater than zero.

For accurate parameter values, the model provides analytical result for real malaria situation. In our case, most of the parameters are from literature hence we can only give a general conclusion about saturation, not necessarily for Western Kenya highlands. This in a way limits the unders-

tanding of the many aspects of this key parameter.

Other important considerations for more comprehensive results would include climatic factors which affect the feeding rate, gonotrophic cycle and mosquito survival.

Our model points to more research, parameter estimation and sensitivity analysis if logical conclusions are to be made.




Figure 5.2 – Variation of susceptible host and vector populations at the Disease Free equilibrium.
 $Q_v = 0.3$ and $\tilde{\mathcal{R}}_0 = 0.718 < 1$




Figure 5.3 – Variation of susceptible host and vector populations at the Endemic equilibrium.
 $Q_v = 0.09$ and $\tilde{\mathcal{R}}_0 = 2.495 > 1$




Figure 5.4 – Variation of Infected Host, Infected Vector, Total Host and Total Vector populations at the Disease Free equilibrium. $Q_v = 0.3$ and $\tilde{\mathcal{R}}_0 = 0.718 < 1$




Figure 5.5 – Variation of Infected Host, Infected Vector, Total Host and Total Vector populations at the Endemic Equilibrium. $Q_v = 0.09$ and $\tilde{\mathcal{R}}_0 = 2.495 > 1$

CONCLUSION AND PERSPECTIVE

We formulated and studied two models for highland malaria transmission in Western Kenya. Some basic models in disease dynamics and epidemiology useful to this thesis have been reviewed.

Our first interest was modeling the discontinuous spatialization, exhibited by the different terrain ecosystems. Considering the importance of the individual's age in the malaria infection and transmission, we grouped the metapopulation in each patch as children (0-5 years) and adults (over 5 years). With a set of differential equations the dynamics of malaria transmission in children and adults in n different patches was described by a system of first order, ordinary differential equations.

A qualitative analysis was done for the age structured-metapopulation model by applying dynamical systems theory. A threshold $\mathcal{R}_0$ was computed, which completely determines if the disease dies out or persists in the population. We prove that, if $\mathcal{R}_0 \leq 1$, the disease free equilibrium is globally stable, and, if $\mathcal{R}_0 > 1$, the endemic equilibrium is unique and globally asymptotically stable. An example of the model on two patches is given. We also provide some numerical study applied to four sites in Western Kenya grouped as U-shaped and V-shaped valley ecosystems. The simulation was done with data from Western Kenya where available and also from published literature. Different scenarios were analysed and the model validated to highlight the importance of metapopulation and age structuring which are often neglected in many malaria models.

In the second model, we studied the dynamics of malaria transmission in the highlands, with disease induced deaths and varying vector and human populations, and saturation in the vector feeding rate. In this model the biting rate is density dependent, and the two population density ratios play a vital role in the transmission. We assume that if the host-vector ratio is low, the bites are few and therefore the probability of transmission reduces too. We introduce a threshold ratio Q_v , such that, as the ratio increases the biting rate, will rise as a function of the density ratio until it reaches this threshold and it becomes a constant. The parameter Q_v is important in our model as it determines which of the two population densities is driving the biting rate hence malaria transmission.

We assume that the human populations vary with time since malaria causes deaths, hence can

have profound effects on the population size. A similar assumption holds for the vector population. A qualitative study of this model has been done and a basic reproduction number calculated. As in the previous case, the existence of equilibrium solutions and their stability depends on the basic reproductive number, $\mathcal{R}_0$ which in turn depends on Q_v . When $\mathcal{R}_0 \leq 1$, we proved that the disease free equilibrium is locally and globally asymptotically stable, and if $\mathcal{R}_0 > 1$, there exists a unique endemic equilibrium that is locally and globally asymptotically stable. The model was validated with some realistic data to show the populations and the role of Q_v on the basic reproduction number, $\mathcal{R}_0$.

Our analysis points to the fact that to fully understand the transmission dynamics of highland malaria in Kenya, and consequently the resurgent epidemics, instead of modeling the factors in isolation, it would be meaningful to characterize fully the the demography and life style of the adult mosquito vector, the age of the hosts, the dynamics of the patches, the seasonality of the epidemics, and the other factors at play in the highlands. This brings out the challenging nature of the malaria control problem and at the same time presents a wonderful opportunity for research and other tools in order to understand what this might mean for for the control of highland malaria in western kenya.

A desirable control policy should be one that target the children population as they are the provide the greatest source of mosquito infection. It should seek to limit contact between mosquitoes and humans and disrupt breeding sites or move them further away from human habitats, so that mosquitoes seeks other feeding sources like animals, which would interrupt the malaria infection. The seasonal of the epidemics in the highland should also guide control policies so that measures are put in place to arrest the spread before the end of the rainy season.

We have presented in this thesis, a framework that researchers can build upon and apply the methodology to other vector-borne diseases , I hope the insights provides information that could potentially lead to eradicating malaria or significantly reducing the disease burden in Western Kenya as it endeavors to achieve the millenium development goals (MDG).

Our attempt towards the search for a practical solution on elimination and control of highland malaria in Western Kenya, is only but a small step in the right direction.

MATHEMATICAL TOOLS

Mathematical modeling of infectious diseases is done by analysing the dynamical systems developed during model formulation. The systems may be ordinary, partial or difference differential equations, which may be linear or nonlinear. In this thesis we have dealt with nonlinear ordinary differential equations and hereby give some definitions and classical results from dynamical systems theory .

A.1 Preliminaries and Notation

Ordered vector space in $\mathbb{R}^n$

We use the standard notation $x \in \mathbb{R}^n$ to define a vector and x_i to denote the i^{th} vector component in $\mathbb{R}^n$. The vector space $\mathbb{R}^n$ is said to be ordered if, when $x \geq y$ the inequality $x_i \geq y_i$ also holds. On the positive orthant the inequality $x \geq y$ is equivalent to $x - y \in \mathbb{R}^n$ and in particular

$$x \geq 0 \Leftrightarrow \text{for every index } i, \quad x_i \geq 0.$$

We write $x > 0$, when $x \geq 0$ and $x \neq 0$, and write $x \gg 0$, if x is in the interior of $\mathbb{R}_+^n$ such that

$$x \gg 0 \Leftrightarrow \text{for every index } i, \quad x_i > 0.$$

The same notation is used for matrices and we define a matrix $A(n, n, \mathbb{R}^{n^2})$ as an $n \times n$ matrix in $\mathbb{R}^{n^2}$. The notation $A \geq B$ holds if for the pair of indices (i, j) we have $a_{ij} \geq b_{ij}$ and $A > B$ holds if $A \gg B$.

For the ordered vector space $\mathbb{R}^n$, we define a closed interval

$$[a, b] = \{x \in \mathbb{R}^n \mid a \leq x \leq b\} = [a_1, b_1] \times \dots \times [a_n, b_n].$$

This notation can also be used on the open interval

$$]]a, b[[= \{x \in \mathbb{R}^n \mid a \ll x \ll b\} =]a_1, b_1[\times \dots \times]a_n, b_n[.$$

If E and F are subsets of $\mathbb{R}^n$, we have the classical definition

$$\mathbb{R}_+ E = \{\lambda x \mid \lambda \in \mathbb{R}_+ \quad x \in E\}$$

and

$$E + F = \{x + y \mid x \in E \ y \in F\}.$$

A scalar product of two vectors is denoted by $\langle x \mid y \rangle$. If M is a matrix and M^T its transpose, we can define the vectors on $\mathbb{R}^n$ and the $n \times 1$ column vectors and express the scalar product as $\langle x \mid y \rangle = x^T y$. We denote by e_i the canonical basis of $\mathbb{R}^n$.

Definition A.1 (Autonomous Systems). Let Ω be a subset of $\mathbb{R}^n$. Consider the autonomous differential equation defined by :

$$\dot{x} = X(x) = f(x) \quad x \in \Omega \quad (\text{A.1})$$

Suppose that $X : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous and satisfies the conditions as a solution of [A.1](#), is unique and continuously depends on the initial conditions. The stationary or equilibrium points of the system [A.1](#) are the points $x_0 \in \Omega$ satisfying $X(x_0) = 0$. For each $x \in \Omega$, we denote by $X_t(x)$ the solution of the system [A.1](#) satisfying $X_0(x) = x$. We suppose that X satisfies the conditions that $X_t(x)$ is continuous in (t, x) .

A.2 Stability

A system which lacks stability would be a poor model for reality as 'reality is always a perturbation of what we think it is'. So some kind of stability is needed in modeling. Two kinds of stability are of great importance in application of differential equations : stability respecting perturbation of initial values for a fixed equation and stability respecting perturbations of the equation itself. In the first case we say the system is 'persistent' and second case 'robust'. We say that an equilibrium point is locally stable if all solutions which start near $\bar{x}$ (meaning that the initial conditions are in the neighborhood of $\bar{x}$) remain near $\bar{x}$ for all future time.

A.2.1 Stability : Basic definition

Consider system [A.1](#).

Definition A.2 (Equilibrium point). A point $\bar{x} \in \mathbb{R}^n$ is an equilibrium point of the system [A.1](#) if $f(\bar{x}, t) = 0$.

Definition A.3 (Lyapunov stability). Let $\bar{x} \in \omega$ be an equilibrium point. System [A.1](#) is stable (we say also Lyapunov stable) at $\bar{x}$ or $\bar{x}$ is a stable equilibrium position for [A.1](#), if for each $\epsilon > 0$ there exists a positive real number δ such that for each x with $|x - \bar{x}| < \delta$, the solution $X(x(t))$ is defined for all $t \geq 0$ and satisfies $|X(x(t)) - \bar{x}| < \epsilon$ for all $t > 0$. When [A.1](#) is not Lyapunov stable at $\bar{x}$, we say that it is unstable at $\bar{x}$.

Definition A.4 (Attractivity). The steady state $\bar{x}$ is said to be attractive (We say also that [A.1](#) is attractive at $\bar{x}$) if there exists a neighborhood $U \in \Omega$ of $\bar{x}$ in such a way that for any initial condition x belonging to U , the corresponding solution $X(t(x))$ of [A.1](#) is defined for all $t \geq 0$ and tends to $\bar{x}$ as t tends to infinity, i.e., $\lim_{t \rightarrow +\infty} X(x(t)) = \bar{x}$.

Definition A.5 (Asymptotic stability). We say that $\bar{x}$ is stable if solutions starting "close" to it at a given time, remain close to it for all future times. It is said to be asymptotically stable if nearby solutions actually converge to $\bar{x}$ at $t \rightarrow \infty$. That means it is Lyapunov stable and attractive.

Definition A.6 (Exponential stability). The system A.1 is exponentially stable (respectively globally exponentially stable) at $\bar{x}$, if there exist two positive constants K and λ such that $|X(x) - \bar{x}| < K|x - \bar{x}|e^{-\lambda t}$ for all x in a neighborhood of $\bar{x}$ (respectively for all $x \in \Omega$) and all positive time t .

Definition A.7 (Attractor). This refers to a compact, nonempty set K which attracts some neighborhood N of itself. It is assumed that K is invariant, that is, it contains the orbits of all its equilibrium points. The neighbourhood N can always be chosen to be invariant also by simply replacing it with the union of all its points. The largest of such N , ie. the set of all points attracted to K is called the basin of K . An attractor enjoys some kind of stability. Any trajectory starting near it may wander away, but eventually returns to approach it asymptotically.

Definition A.8 (Global stability). We say an equilibrium point $\bar{x}$ is globally stable if it is stable for all initial conditions $x_0 \in \mathbb{R}^n$

A.2.2 Dynamical Properties

Definition A.9 (Invariant set). Given the autonomous dynamical system $\dot{x} = f(x)$ and a trajectory $s(t, x_0)$ where x_0 is the initial point. Let

$$\mathcal{O} \triangleq \{x \in \mathbb{R}^n \mid \phi(x) = 0\}$$

where ϕ is a real valued function. Then the set $\mathcal{O}$ is said to be positively invariant if $x_0 \in \mathcal{O}$ implies that $x(t, x_0) \in \mathcal{O}$, for all $t \geq 0$. This means that once a trajectory of the system enters $\mathcal{O}$, it will not leave it again.

Definition A.10 (Orbit). The orbit $\gamma^+(x_0)$ is called a positive orbit if for all x_0 in the set

$$\{x(t, x_0) \mid t \geq 0\},$$

the orbit is defined by :

$$\gamma(x_0) = \{x(t, x_0) \mid t \in \mathbb{R}\}.$$

The set is positively invariant if $\gamma^+(M) \subset M$, and invariant if it contains the orbits of each of its points.

Note : In the study of dynamical systems, a limit set is the state a dynamical system reaches after an infinite amount of time has passed, by either going forward or backwards in time. Limit sets are important because they can be used to understand the long term behavior of a dynamical system.

Definition A.11 (ω -Limit point). A point p is called an ω -limit point of $X_t(x)$ if there exists a sequence $t_n \in \mathbb{R}$ such that

$$\lim_{n \rightarrow +\infty} t_n = +\infty \quad \text{and} \quad \lim_{n \rightarrow +\infty} x_{t_n}(x) = p$$

The set of all ω -limit point is the ω -limit set of x and is denoted by $\omega(x)$. This means that the sequence t_n tends to $+\infty$ as n tends to infinity and the flow through x tends to p as n tend to $+\infty$

Theorem A.1. If the positive orbit $\gamma^+(x_0)$ is bounded then the set of points ω -limit points, $\omega(\gamma)$ is non-empty, compact, connected and invariant.

Theorem A.2 (Poincare-Bendixon). Consider the equation $\dot{x} = f(x)$ in $\mathbb{R}^2$. Suppose that γ^+ is a bounded positive orbit and $\omega(\gamma^+)$ does not contain equilibrium points, then $\omega(\gamma^+)$ is a periodic orbit. If $\omega(\gamma^+) \neq \gamma^+$ this periodic orbit is called a limit cycle.

Definition A.12. For the $\mathcal{C}^1$ autonomous system $\dot{x} = f(x)$ and an equilibrium point x_0 , the linearised system in x_0 is defined by the linear system

$$\dot{x} = Df(x_0)x,$$

where $Df(x_0)$ is the derivative of f at x_0 .

Theorem A.3 (Poincare-Lyapunov). Consider the $\mathcal{C}^1$ system $\dot{x} = f(x)$ and an equilibrium point x_0 .

1. If $Df(x_0)$ has the real parts of all its eigenvalues negative, then x_0 is asymptotically stable.
2. If $Df(x_0)$ has (at least) one of its eigenvalues with a positive real part, then x_0 is unstable.

Theorem A.4 (Barrier Theorem [21, 127]). Consider a function $H : \mathbb{R}^n \rightarrow \mathbb{R}$ and an open set Ω . We assume that in every point $x \in \bar{\Omega}$ such that $H(x) = 0$ we have $\nabla H(x) \neq 0$ and

$$X(x) \mid \nabla H(x) \neq 0, x \in \bar{\Omega} \leq 0$$

Then for every trajectory of X starting from a point satisfying $H(x) \leq 0$ cannot get from the set

$$x \in \mathbb{R}^n \mid H(x) = 0, x \in \bar{\Omega} \leq 0$$

Proof




Figure A.1 – Vector fields pointing toward the interior

The boundary of the nonnegative orthant, is the reunion of the hyperplane

$$H_i = \{x | x_i = 0\}$$

Let $x \in H_i, x \geq 0$, then

$$\dot{x}_i = (Ax)_i = \sum_{j=1}^n a_{i,j}x_j = a_{i,i}x_i + \sum_{j \neq i}^n a_{i,j}x_j = \sum_{j \neq i}^n a_{i,j}x_j \geq 0$$

The vector fields points toward the interior and hence Barrier theorem shows invariance

A.3 Monotone systems

Consider the system A.1 where X is $\mathcal{C}^1$ and Ω is an open set in $\mathbb{R}^n$.

- X is said to be of type K in Ω if for each i ; $X_i(a) \leq X_i(b)$ for any two points a and b in Ω satisfying $a_k \leq b_k$ and $a_i = b_i$, $(i \neq k \text{ and } i, k = 1, 2, \dots, n)$;
- We say that Ω is p -convex if $tx + (1-t)y \in \Omega$, for all $t \in [0, 1]$ whenever $x, y \in \Omega$ and $x \leq y$;
- The system A.1 is said to be a cooperative system if Ω is P -convex and

$$\frac{\partial X_i(x)}{\partial x_j} \geq 0, \quad i \neq j, \quad x \in \Omega$$

- We say that system A.1 is a competitive system if Ω is P -convex and

$$\frac{\partial X_i(x)}{\partial x_j} \leq 0, \quad i \neq j, \quad x \in \Omega$$

A.3.1 Monotone Dynamical System

Consider a dynamical system [72] with a flow $\phi_t : x \rightarrow \phi_t(x)$. This dynamical system is said to be monotone if it is defined on an ordered metric space with the following property ;

$$t \geq 0, \quad x \leq y \Rightarrow \phi_t(x) \leq \phi_t(y)$$

It is said to be strongly monotone if

$$t \geq 0, \quad x < y \Rightarrow \phi_t(x) \ll \phi_t(y)$$

We say that the system is anti-monotone if

$$t \geq 0, \quad x \leq y \Rightarrow DX(x) > DX(y), \text{ and}$$

It is strictly anti-monotone if

$$t \geq 0, \quad x < y \Rightarrow DX(x) > DX(y).$$

A.3.2 Triangular System

A triangular system [158] is precisely an $\mathbb{R}^n \times \mathbb{R}^m$ system of the form

$$\begin{cases} \dot{x}_1 = f_1(x_1) \\ \dot{x}_2 = f_2(x_1, x_2) \end{cases} \quad (\text{A.2})$$

where f_1 is a map from $\mathbb{R}^n$ to $\mathbb{R}^n$ and f_2 from $\mathbb{R}^n$ to $\mathbb{R}^m$. We suppose that the conditions for existence and uniqueness of solutions are satisfied, (eg. f_1 and f_2 are $\mathcal{C}^1$). The trajectories of the system have the same projection on $\mathbb{R}^n \times \{0\}$ and hence the name triangular. In fact the Jacobian of this system is a lower triangular block, and it is also said to be hierarchical. We will deduce a stability result from Vidyasagar Theorem, and present a different and simpler version than the one given by Vidyasagar.

Theorem A.5 (Vidyasagar). Consider the following $\mathcal{C}^1$ system :

$$\begin{cases} \dot{x}_1 = f_1(x_1) \\ \dot{x}_2 = f_2(x_1, x_2) \end{cases} \quad (\text{A.3})$$

If the origin of $\mathbb{R}^n$ is globally asymptotically stable (GAS) for the system $\dot{x}_1 = f_1(x_1)$ in $\mathbb{R}^n$ and the origin of $\mathbb{R}^m$ is GAS for $\dot{x}_2 = f_2(0, x_2)$ on $\mathbb{R}^m$, then the origin of $\mathbb{R}^n \times \mathbb{R}^m$ is asymptotically stable. Furthermore if all the trajectories are bounded, then the origin is GAS for A.3 on $\mathbb{R}^n \times \mathbb{R}^m$.

Proof of Stability

Let the following be a neighborhood of the the origin

$$B(0, \varepsilon) = \{(x_1, x_2) \mid \|x_2\| \leq \varepsilon\}.$$

Since the equilibrium points of the isolated $\mathcal{C}^1$ system are GAS, we can apply the inverse Lyapunov theorem. There exists positive definite $\mathcal{C}^1$ functions $V_1(x_1)$ and $V_2(x_2)$ such that

$$\dot{V}_1 = \langle \nabla V_1(x_1) \mid f_1(x_1) \rangle \leq 0, \quad \dot{V}_2 = \langle \nabla V_2(x_2) \mid f_2(0, x_2) \rangle \leq 0$$

The functions $\dot{V}_1$ and $\dot{V}_2$ are negative definite on $B(0, \varepsilon)$ for ε sufficiently small since f_1 and V_1 are $\mathcal{C}^1$. Let

$$L = \max_{(x_1, x_2) \in B(0, \varepsilon)} \frac{\partial f_1}{\partial x_2}(x_1, x_2)$$

$$M = \max_{(x_1, x_2) \in B(0, \varepsilon)} \nabla V_2(x_2)$$

Since V_2 is a Lyapunov function, we can choose $\delta_1 < \frac{\varepsilon}{2}$ sufficiently small such that

$$\max_{\|x_2\| \leq \delta_1} \nabla V_2(x_2) < \min_{\frac{\varepsilon}{2} \leq \|x_2\| \leq \varepsilon} \nabla V_2(x_2)$$

$$\dot{V}_2(x_2) = \langle \nabla V_2(x_2) \mid f_2(x_1, x_2) \rangle = \langle \nabla V_2(x_2) \mid f_2(0, x_2) \rangle + \langle \nabla V_2(x_2) \mid f_2(x_1, x_2) - f_2(0, x_2) \rangle$$

From Taylors formular we have the following relation

$$f_2(x_1, x_2) - f(0, x_2) = \int_0^1 \frac{\partial f_2}{\partial x_1}(x_1, x_2) x_1 dt$$

and on $B(0, \varepsilon)$ we have

$$\|f_2(x_1, x_2) - f_2(0, x_2)\| \leq \|x_1\|$$

and the Cauchy Schwarz inequality gives

$$\dot{V}_2(x_2) \leq \langle \nabla V_2(x_2) \mid f_2(0, x_2) \rangle + LM\|x_1\| \quad (\text{A.4})$$

Since the function $\langle \nabla V_2(x_2) \mid f_2(0, x_2) \rangle$ is negative, we can define a function φ by

$$\varphi(c) = \min_{c \leq \|x_2\| \leq \varepsilon} -\langle \nabla V_2(x_2) \mid f_2(0, x_2) \rangle$$

This function φ , defined on $\mathbb{R}$, is continous, decreasing and tend to 0 as $c \rightarrow 0$. We show that $\varphi(c) > 0$ for all $c > 0$.

Since the system $\dot{x}_1 = f_1(x_1)$ is asymptotically stable, we can choose $\delta_2 \leq \delta_1$ such that for all initial conditions satisfying $\|x_1(0)\| \leq \delta_2$, and $t \geq 0$, we have the inequality $\|x_1(t)\| \leq \frac{\varphi(c)}{LM}$. Moreover if we have $\|x_1\| \leq \delta_2$ and $\|x_2\| \geq \delta_1$, with the inequality A.4, we deduce the relation

$$\langle \nabla V_2(x_2) \mid f_2(0, x_2) \rangle + LM\|x_1\| < 0 \quad (\text{A.5})$$

We define $0 < \delta_3 < \delta_2$ so that

$$\max_{\|x_1\| \leq \delta_3} V_1(x_1) < \min_{\delta_2 \leq \|x_1\| \leq \varepsilon} V_1(x_1)$$

Consider the open set $\mathcal{U}$ defined by

$$\mathcal{U} = \{(x_1, x_2) \mid \|x_1\| \leq \delta_2, \|x_2\| \leq \delta_3\}.$$

If $x_1(0) \leq \delta_3$, since V_1 is decreasing, the preceding inequality implies that $\|x_1(t)\| \leq \delta_2$ (all the trajectory are inside the sphere of radius δ_2 in $\mathbb{R}^n$).

For $x_2(0) \leq \delta_3$, we have

$$\max_{\|x_2\| \leq \delta_3} V_2(x_2) \leq \max_{\|x_2\| \leq \delta_3} \|x_2\| \leq \delta_1 V_2(x_2) < \min_{\frac{\varepsilon}{2} \leq \|x_2\| \leq \varepsilon} V_2(x_2).$$

The trajectory $(x_1(0), x_2(0))$ satisfy $\|x_2(t)\| \leq \delta_1$ which implies

$$V_2(x_2) \leq \min_{\delta_1 \leq \|x_2\| \leq \varepsilon} V_2(x_2).$$

Consequently, we have $\|x_2\| \leq \delta_2$. This leads to the inequality $\|x_2(t)\| \geq \delta_1$, and from the inequality A.5, we have $\dot{V}_2 \leq 0$.

Since V_2 is decreasing on the trajectories contained in the ring $\|x_1\| \leq \delta_2, \|x_2\| \leq \frac{\varepsilon}{2}$, we note that a trajectory cannot reach the sphere of radius $\frac{\varepsilon}{2} \in \mathbb{R}^m$. This shows that $\|x_1(t)\| \leq \delta_2 \leq \varepsilon$ and $\|x_2(t)\| \leq \frac{\varepsilon}{2}$, which completes the prove of stability.

We will now show the local attractivity by the Lasalle invariance principle.

Since the origin is stable, there exists a compact set X and a neighborhood U of the origin which is positively invariant. This restricts us to the invariant set U .

Consider the Lyapunov-Lasalle function V_1 . By the hypothesis

$$\dot{V}_1 = \langle \nabla V_1(x_1) \mid f_1(x_1) \rangle \leq 0.$$

Let

$$E = \{(x_1, x_2) \in \mathcal{U} \mid \dot{V}_1(x) = 0\}$$

We also consider the largest invariant set E . This is evidently the set $\{0\} \times \mathbb{R}^m \cup \mathcal{U}$. By hypothesis the system $\dot{x}_2 = f_2(0, x_2)$ is globally asymptotically stable in $\{0\} \times \mathbb{R}^m$. This implies that any negative trajectory of the system $\mathcal{U} \setminus \{0\}$ exit $\mathcal{U}$.

In effect if that is not the case, then there exists a trajectory $\gamma \in \mathcal{U}$. Then the set of α -limit points of γ are invariant. By the asymptotic stability and invariance, this point contains the origin. This means that the trajectory starting close to the origin remain close to it. The fact that this trajectory is invariant contradicts the stability so the property stated is true. This means that the largest invariant set containing E is reduced to the origin. Which shows that $\mathcal{U}$ is attractive.

If a trajectory is relatively compact, then the ω -limit points are in $\{0\} \times \mathbb{R}^m$. In effect, for $t_n \rightarrow \infty$, we have $x_1(t_n) \rightarrow 0$. If all the trajectories are compact, then the ω -limit points are in $\{0\} \times \mathbb{R}^m$. By the asymptotic stability of $\{0\} \times \mathbb{R}^m$, the origin is an ω -limit. All trajectories get as close as possible to the origin. Due to stability they are attracted to the open set $\mathcal{U}$ defined above, otherwise they approach the origin asymptotically.

A.3.3 Lyapunov Methods

The Lyapunov function plays a major role in the study of the stability of dynamical systems. We dedicate this section to some results due to Lyapunov. Let $V : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function.

Definition A.13. We have the following definitions

1. The function V is said to be positive definite if $V(x) = 0$ and $V(x) > 0$ in a neighborhood Ω_0 of x_0 for all $x \neq x_0$ in the neighborhood.
2. The function V is said to be negative definite if $-V$ is positive definite.
3. The function V is said to be semi-positive if $V(x_0) = 0$ and $V(x) \geq 0$ in a neighborhood Ω of x_0

Theorem A.6 (Lyapunov Theorem). Let V be a function

- If a function V is positive definite and $\dot{V}$ is negative semi-definite in Ω , then the equilibrium point x_0 is stable for the system [A.1](#).
- If the function V is positive definite and $\dot{V}$ is negative definite in Ω , then the equilibrium point x_0 is asymptotically stable for the system [A.1](#)

This theorem implies that to show that an equilibrium point x_0 is stable, it is sufficient to find a Lyapunov function for the point. Moreover, to use the original Lyapunov theorem to show the asymptotic stability of a given system, we must determine a function V whose derivative is positive definite and the derivative $\dot{V}$ is negative definite. In a general case, this is not straightforward. The condition on the derivative $\dot{V}$ can be relaxed by using the LaSalle Invariance principle introduced in the next section.

A.3.4 LaSalle Invariance Principle

Theorem A.7 (LaSalle Invariance Principle [98, 99]). Let $\Omega \subset \mathbb{R}^n$ be a compact set that is positively invariant with respect to the system A.1. Let $V : \Omega \rightarrow \mathbb{R}$ be continuously differentiable such that $\dot{V}(x) \leq 0$ in Ω . Let E be the set of all points in Ω where $\dot{V}(x) = 0$. Let L be the largest invariant set in E . Then every solution starting in Ω approaches L as $t \rightarrow \infty$.

This theorem is a very important tool for systems analysis, and is different from Lyapunov, as it does not require V to be positive definite and $\dot{V}$ to be negative definite. However, it only provides information on the attractiveness of the considered system at the equilibrium x_0 . For example, it can be used to prove that the solutions tend toward an equilibrium point when the set L is reduced to that equilibrium point. It does not indicate whether this equilibrium is stable or not. When one wants to establish asymptotic stability of an equilibrium $x_0 \in \Omega$, we use the following corollary which is a consequence of the LaSalle invariance principle.

Corollary A.1 (LaSalle,[99]). Consider the compact set $\Omega \subset \mathbb{R}^n$ with $x_0 \in \Omega$. Let $V : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable positive definite function such that $\dot{V} \leq 0$ in Ω . Let $E = \{x \in \Omega \mid \dot{V} = 0\}$. Assume that the largest positively invariant set contained in E is reduced to the point x_0 . Then x_0 is an asymptotically stable equilibrium point for the system A.1. If these conditions are satisfied for $\Omega = \mathbb{R}^n$, if in addition V is in Ω ie. $\lim_{\|x\| \rightarrow \infty} V(x) = +\infty$ when $d(x, \frac{\partial}{\partial x} \Omega) \rightarrow +\infty$, then all trajectories are bounded for $t \geq 0$ and x_0 is a globally stable equilibrium point for the system A.1.

Corollary A.2. Under the assumptions of the previous theorem, if the set L is reduced to the point $x_0 \in \Omega$, then x_0 is a globally stable equilibrium point for the system A.1 defined on Ω .

A.4 Matrices

Definition A.14 (Stability Modulus, Spectral radius). Let A be a square matrix, we denote by $\text{spec}(A)$ all the eigenvalues of A . The stability modulus of A is the number defined by

$$\alpha(A) = \max\{\text{Re}(\lambda); \lambda \in \text{Spec}(A)\}.$$

The matrix A is said to be stable if $\alpha(A) < 0$.

The spectral radius is the real number $\rho(A)$ defined by

$$\rho(A) = \max_{\lambda \in \text{Spec}(A)} |\lambda|.$$

We say that a matrix A is stable if its eigenvalues have strictly negative real parts. Such a matrix is also said to be Hurwitz [143, 149, 151].

Theorem A.8 (Gershgorin circle theorem). Let $A \in \mathcal{M}_n(\mathbb{R})$. The spectrum of A is contained in the union of disks whose centers are the diagonal coefficients a_{ii} of the matrix A and radii are the respective sums of the absolute values of the remaining coefficients in the row.

In other words for each index i , $1 \leq i \leq n$, if we set $r_i = \sum_{j=1, j \neq i}^n |a_{ij}|$, then

$$\text{Spec}(A) \subset \bigcup_{i=1}^n B(a_{ii}, r_i)$$

Lemma 1. Let $A \in \mathcal{M}_n(\mathbb{R})$ be a square matrix. If A is a singular matrix, then there exists an index i_0 such that

$$|a_{i_0 i_0}| \leq \sum_{j=1, j \neq i_0}^n |a_{i_0 j}|.$$

In other words, the term $a_{i_0 i_0}$ dominates the column i_0 .

Proof

Let n be an arbitrary integer. Assume that A is a singular matrix. Then, the matrix A will be the linear map of $\mathbb{R}^n$ to itself, or its corresponding matrix. Since A is singular, $\text{Ker}(A) \neq \{0\}$; Let $x = (x_1, x_2, \dots, x_n)^T \in \text{Ker}(A)$; Let i_0 be an index of the component x such that x_{i_0} achieves the maximum of x (that is to say $|x| = \max_{1 \leq i \leq n} |x_i| = \|x\|_\infty$

$$Ax = 0 \Rightarrow (Ax)_{i_0} = \sum_{j=1}^n a_{i_0 j} x_j = 0$$

$$\Rightarrow -a_{i_0 i_0} x_{i_0} = \sum_{j=1, j \neq i_0}^n a_{i_0 j} x_j = 0$$

$$\Rightarrow |a_{i_0 i_0}| |x_{i_0}| \leq \sum_{j=1, j \neq i_0}^n |a_{i_0 j}| |x_j| = 0$$

$$\leq \sum_{j=1, j \neq i_0}^n |a_{i_0 j}| |x_j| = 0.$$

Since $x_{i_0} \neq 0$, it follows that

$$|a_{i_0 i_0}| \leq \sum_{j=1, j \neq i_0}^n |a_{i_0 j}|$$

which ends the proof of the lemma. The following is a better consequence of the theorem and the preceding lemma;

Corollary A.3. Let $A \in \mathcal{M}_n(\mathbb{R})$ be a square matrix. If A is a singular matrix, then there exists an index i_0 such that

$$|a_{i_0 i_0}| \leq \sum_{j=1, j \neq i_0}^n |a_{i_0 j}|.$$

In other words, the term $a_{i_0 i_0}$ dominates the column i_0 . The proof of this corollary is identical to that of the previous lemma, however we replace the matrix A by its transpose A^T which has the same properties as the matrix A .

Proof of theorem A.8 Let $A \in \mathcal{M}_n(\mathbb{R})$; then for all $\lambda \in \text{Spec}A$, $(A - \lambda I)$ is a singular matrix. The previous lemma implies that for all $\lambda \in \text{Spec}A$, there exists $i_\lambda \in \mathbb{N}$, $1 \leq i_\lambda \leq n$ such that

$$|a_{i_\lambda i_\lambda}| \leq \sum_{j=1, j \neq i_\lambda}^n |a_{i_\lambda j}|$$

If we set $r_i = \sum_{j=1, j \neq i}^n |a_{ij}|$; the previous inequalities are equivalent to; for all $\lambda \in \text{Spec}(A)$, there exists $i_\lambda \in \mathbb{N}$, $1 \leq i_\lambda \leq n$ such that $B(a_{i_\lambda}, r_{i_\lambda})$. If we consider the balls $B(a_{ii}, r_i)$ for all i , $1 \leq i \leq n$, it follows that any $\lambda \in \text{Spec}(A)$ is inside one of them. So we can say that $\lambda \in \cup_{i=1}^n B(a_{ii}, r_i)$. This completes the prove.

Corollary A.4. Let $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ be a square matrix. The spectrum of the matrix A is contained in the union of the disks whose centers are the diagonal coefficients a_{ii} diagonal of the matrix A and whose radii are the sums of the absolute values of the off-diagonal coefficients of the corresponding columns. In other words, for all i , $1 \leq i \leq n$, if we set $r_i = \sum_{j=1, j \neq i}^n |a_{ji}|$, then

$$\text{Spec}(A) \subset \cup_{i=1}^n B(a_{ii}, r_i)$$

Proof

This is a consequence of Theorem A.8 and Corollary A.3. In effect A is strictly diagonally dominant, then for all i such that $1 \leq i \leq n$, we have $|a_{ii}| > r_i$ ($r_i = \sum_{j=1, j \neq i}^n |a_{ji}|$, for the respective columns, or $r_i = \sum_{j=1, j \neq i}^n |a_{ij}|$ for the respective row). So none of the balls $B(a_{ii}, r_i)$ contains the origin of $\mathbb{C}$. Therefore no eigenvalue of A can be zero.

Definition A.15. We have the following definitions

- A matrix $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ is a strictly column diagonally dominant if for all i such that $1 \leq i \leq n$, we have

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ji}|$$

- A matrix $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ is a strictly row diagonally dominant if for all i such that $1 \leq i \leq n$, we have

$$|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|$$

- We say that a matrix is strictly column (respectively row) diagonally dominant if the above inequalities strictly hold.

Corollary A.5. If a matrix A is strictly diagonally dominant, then it is invertible.

This is an immediate consequence of A.8 (Proof by contradiction).

Definition A.16. Let $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ be a square matrix. We say that the matrix A is reducible, if there exists a permutation matrix P such that

$$P^T A P = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix} \quad (\text{A.6})$$

The matrix A is said to be irreducible if and only if it is not reducible.

A.4.1 Metzler Matrices

Definition A.17. We say that $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ is a Metzler matrix if all the off-diagonal entries are non-negative. That is, $a_{ij} \geq 0$ for all i and j with $i \neq j$ [18].

The following theorem will be very useful for studying the stability of our models.

Theorem A.9 (Perron-Frobenius Theorem, [18]). If $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ is a non negative matrix ;

- Then the spectral radius $\rho(A)$ is an eigenvalue of A and there is a corresponding eigenvector $v > 0$
- If in addition the matrix A is irreducible, then $\rho(A) > 0$ and $v \gg 0$; moreover, $\rho(A)$ has algebraic multiplicity 1 and if $u > 0$ is an eigenvector of A , then there exists $s > 0$ such that $u = s v$.
- If $B > A$, then $\rho(B) > \rho(A)$.
- Finally, if $A \gg 0$, then $|\lambda| < \rho(A)$ for all other eigenvalues λ of A .

Corollary A.6. Let $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ be a Metzler matrix.

- The stability modulus $\alpha(A)$ is an eigenvalue of A and there exists a vector $v > 0$ such that $Av = \alpha(A)v$. Moreover $Re(\lambda) < \alpha(A)$ for all $\lambda \in Spec(A) - \{\alpha(A)\}$.
- If in addition A is irreducible, then
 - (a) $\alpha(A)$ is a simple eigenvalue;
 - (b) $v \gg 0$ and all other positive eigenvectors of A are multiples of v ;
 - (c) If B satisfies $B > A$, then $\alpha(B) > \alpha(A)$;
 - (d) If $\alpha(A) < 0$, then $-A^{-1} \gg 0$.

Proof

Let $C \geq 0$ be real and large enough such that $A + CI \geq 0$. Hence applying the Perron-Frobenius theorem defined above on $A + CI$ implies that $C \geq 0$. In particular, the spectral radius $\rho(A + CI) > 0$. We note that $Spec(A + CI) = C + Spec(A)$; which implies that

$$\alpha(A + CI) = \rho(A + CI) = C + \alpha(A)$$

and consequently, $\alpha(A)$ is an eigenvalue of A . If A is irreducible, $A + CI$ is also irreducible.

The theorem of Perron-Frobenius theorem applied to A implies properties 1, 2, 3. Moreover as $-a^{-1} = \int_0^\infty e^{tA} dt \gg 0$, (4) is satisfied. The following theorem is important to study the stability of equilibrium points and the construction of Lyapunov functions.

Theorem A.10 (Metzler Stable Matrices). Let $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ be a Metzler matrix, the following statements are equivalent :

- The Matrix A is Metzler stable,
- The Matrix A is invertible and $-A^{-1} \geq 0$,
- If $b \gg 0$, then the equation $Ax + b = 0$ has a solution $x \gg 0$,
- There exists a vector $c > 0$, such that $Ac \ll 0$,
- There exists a vector $c \gg 0$, such that $Ac \ll 0$

The proof given here is an adaptation of a similar proof given in [18].

Proof :

1) $\Rightarrow$ 2)

The matrix A is stable, invertible and $\alpha(A) < 0$; there exists a scalar k such that for all

$$x_0 \in \mathbb{R}_n, \|e^{tA} x_0\| \leq K e^{\alpha(A)t} \|x_0\|.$$

We deduce that the integral $\int_0^\infty e^{tA} x_0 dt$ is normally convergent for all x_0 , therefore the matrix

$\int_0^\infty e^{tA} dt$ is absolutely convergent

In effect the function $t \in \mathbb{R}_+ \rightarrow e^{tA} \in \mathcal{M}_n(\mathbb{R})$ is in $L^1(0, \infty)$ and

$$B = \int_0^\infty e^{tA} dt = -A^{-1}$$

We will determine the sign of each of the coefficients of the matrix $B = (b_{ij})$; For this we consider the canonical base (e_i) in $\mathbb{R}^n$; we note that;

$$\begin{aligned} b_{ij} &= \langle B e_j, e_i \rangle = (B e_j)_i = \left\langle \int_0^\infty e^{tA} dt e_j, e_i \right\rangle \\ &= \int_0^\infty \langle e^{tA}, e_i \rangle dt \\ &= \int_0^\infty (e^{tA} e_j)_i dt \geq 0 \end{aligned}$$

Hence the matrix $-A^{-1} = B \geq 0$.

2) $\Rightarrow$ 3)

Let b be a vector such that $b \gg 0$, and $x = -A^{-1}b$; then $x \gg 0$ is a product of two positive matrices. The matrix $Ax + b = 0$

3) $\Rightarrow$ 4)

Let $b \gg 0$ be a vector in $\mathbb{R}_+^n$; Let $c = -A^{-1}b$; it is clear that $Ac = -b \ll 0$.

4) $\Rightarrow$ 5)

Perturbing property (4) a little, we set $\varepsilon > 0$ and $c_1 = c + \varepsilon \sum_{i=1}^n e_i \gg 0$.

So $A c_1 = A c + \varepsilon \sum_{i=1}^n A e_i$. By continuity we can choose $\varepsilon > 0$ sufficiently small such that $A c_1 \ll 0$.

5) $\Rightarrow$ 1)

Consider the differential equation $\dot{x} = A^T x$ in the positive orthant. We choose a function

$$V(x) = \langle c, x \rangle.$$

From the hypothesis $c \gg 0$, V is positive definite in $\mathbb{R}_+^n$ and so

$$\dot{V}(x) = \langle c, Ax \rangle = \langle A^T c, x \rangle.$$

$\dot{V}(x) = 0$, if and only if $x = 0$. The Lyapunov theorem proves that the matrix A^T is asymptotically stable, hence A is stable since A and A^T have the same eigenvalues. This completes the prove of the theorem.

Theorem A.11. Let $A = (a_{ij}) \in \mathcal{M}_n(\mathbb{R})$ be a Metzler matrix.

1. The stability modulus $\alpha(A)$ of A is an eigenvalue of A with an associated positive eigenvector $v \in \mathbb{R}_+^n$ such that $v \neq 0$ and $Av = \alpha(A)v$;
2. If in addition the matrix A is irreducible, then $\alpha(A)$ is a simple eigenvalue of the matrix A to which a positive eigenvector is associated. That is to say, there exists $v \in \mathbb{R}_+^n$ such that $v \gg 0$ and $Av = \alpha(A)v$.

Proof

Let $m = \min\{\min_{1 \leq i \leq n} a_{ii}, 0\}$ then $A - mI \geq 0$. Applying the Perron-Frobenius theorem on the matrix $A - mI$ it follows that there exist

$$v \in \mathbb{R}_+^n; (A - mI)v = \rho(A - mI)v$$

or there exists $v \in \mathbb{R}_+^n$ such that

$$Av = (\rho(A - mI) + m)v$$

the vector v is an eigenvector of the matrix A associated with the eigenvalue $\rho(A - mI) + m$.

We note that for a square matrix B we have

$$\text{Spec}(B + kI) = k + \text{Spec}(B)$$

or $\rho(A - mI) = \max_{\lambda \in \text{Spec}(A - mI)} \text{Re}(\lambda)$ or

$$(\rho(A - mI) + m) = \max_{\lambda \in \text{Spec}(A - mI + mI)} \text{Re}(\lambda) = \max_{\lambda \in \text{Spec}(A)} \text{Re}(\lambda) = \alpha(A).$$

So v is an eigenvector of A associated with the eigenvalue

$$\alpha(A) = \rho(A - mI) + m \in \mathbb{R}.$$

Although this proof can be found in [132], we have given the complete proof because the theorem is very crucial in this thesis.

A.4.2 Characteristics of Metzler stable matrices

Let M be a Metzler matrix of the form

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$$

M is stable if and only if A and $D - CA^{-1}B$ are stable.

Proof

We first prove the necessary condition. Since any principal sub-matrix of a Metzler stable matrix is still a Metzler stable matrix, (see [152]), then A and D are Metzler stable. As A is Metzler stable, there exists a vector $c = (c_1, c_2) \gg 0$, such that $Mc \ll 0$ (condition I_{28} of theorem 2.3 [18]). This means that

$$A c_1 + B c_2 \ll 0,$$

$$C c_1 + D c_2 \ll 0.$$

Since A is Metzler stable, $-A^{-1} \geq 0$ and C is positive. Multiplying $A c_1 + B c_2 \ll 0$ by $-C A^{-1}$, we obtain $-C C_1 - C A^{-1} B c_2 \ll 0$ and consequently $(D - C A^{-1} B) c_2 \ll 0$. This proves, from I_{28} that $(D - C A^{-1} B)$ is Metzler stable. This is a necessary condition.

We now prove the sufficient condition.

Since A and $(D - C A^{-1} B)$ are Metzler stable, there exists a vector $c_2 \ll 0$ such that

$$(D - C A^{-1} B) c_2 \ll 0.$$

So we now have $c_3 = -A^{-1} B c_2$, since A is Metzler stable, we have $-A^{-1} \geq 0$. As B is positive and $c_2 \gg 0$, we deduce that $c_3 \gg 0$ and the inequalities

$$C c_3 + D c_2 \ll 0, \quad A c_3 + B c_2 \ll 0.$$

A is Metzler stable, so there exists a vector $v \gg 0$ such that $A v \ll 0$ (see theorem A.10). Now we choose $c_1 = c_3 + \varepsilon v \gg 0$, for $\varepsilon > 0$. We have

$$C c_1 + D c_2 = C c_3 + D c_2 + \varepsilon C v.$$

One can choose $\varepsilon > 0$ sufficiently small so that $C c_1 + D c_2 \ll 0$. Now,

$$A c_1 + B c_2 = A c_3 + \varepsilon A v \ll 0.$$

So we find that $c = (c_1, c_2) \gg 0$ such that $Mc \ll 0$. which proves that M is Metzler stable (see theorem A.10). This proves the sufficient condition. In the following result we will show that the stability modulus is an increasing function of Metzler matrices.

Theorem A.12 (Monotonicity and Stability Modulus [72]).

1. If there exist $v \gg 0$ such that $Av \leq \beta v$, then $\alpha(A) \leq \beta$.
2. If in addition, A is irreducible then $v > 0$ and $Av < \beta v$ leading to $\alpha(A) < \beta$. Infact we must have $v \gg 0$.
3. If there exists $v > 0$ such that $sv \leq Av$, then $s \leq \alpha(A)$.
4. If in addition A is irreducible, then $s < Av$ implies $s < \alpha(A)$

A.4.3 Irreducible Metzler Matrices

In this section we will characterize a dynamical property of irreducible Metzler matrices. A Metzler matrix is invariant on the positive orthant. First, we prove a result on the irreducibility. The definition of irreducibility for Metzler matrices is equivalent to the following property,

Proposition 3 (Irreducible Metzler Matrices). A Metzler matrix A is irreducible if and only if, for any vector $x > 0$ belonging to a face F of $\mathbb{R}_+^n$, where F is defined by :

$$F = \{x \geq 0 \mid \exists I \in I, \langle e_i \mid x \rangle = 0\}$$

there exists an index i such that $\langle e_i \mid x \rangle = 0$ and $\langle e_i \mid Ax \rangle > 0$.

Proof

From theorem A.13, the irreducibility of a matrix depends only on its off-diagonal terms. We can always replace A with $A + \lambda I$ for λ sufficiently large. The irreducibility condition is equivalent to the proposition condition. If there exists i such that $\langle e_i \mid x \rangle = 0$ and $\langle e_i \mid Ax \rangle > 0$, then this is equivalent to $\langle e_i \mid x \rangle = 0$ and $\langle e_i \mid (A + \lambda I)x \rangle > 0$. Without loss of generality we will therefore assume that $A \geq 0$.

The sufficient condition.

As a consequence, we suppose that for all i such that $\langle e_i \mid x \rangle = 0$, we have $\langle e_i \mid Ax \rangle > 0$. Let $F_x = \mathbb{R}^+[0, x]$ be the face generated by x . Since $A \geq 0$, we have $AF_x = \mathbb{R}^+[0, Ax]$. The face F_x is characterized by the set of indices I . Now we have $F_x = \{x \geq 0 \mid \langle e_i \mid x \rangle = 0\}$. For these indices we have $\langle e_i \mid x \rangle = 0$. As a result $AF_x \subset F_x$ and the matrix A is not irreducible.

The necessary condition.

If A is reducible, there exists a face denoted by F_x , such that, $AF_x \subset F_x$. For any index i such that $\langle e_i \mid x \rangle = 0$ we have $\langle e_i \mid Ax \rangle = 0$.

Definition A.18. A directed graph $\mathcal{G} = (X, U)$ is a pair consisting of a set $X = \{x_1, \dots, x_n\}$ and a set U of $X \times X$. The elements of X are called nodes or vertices of the graph. An element $(x, y) \in U$ is called an arc, x is the origin and y is the endpoint. A directed graph (resp. undirected) is a set of vertices with directed arcs (resp. undirected) linking the vertices.

Definition A.19. A path is a sequence of arcs $(u_1, \dots, u_p)$ such that each arc u_i has the endpoint which is the origin of u_{i+1} . It is said that the path joins the origin u_1 with the end u_p . A graph is strongly connected if every pair of distinct vertices is joined by a path.

Any square matrix is associated with a directed graph.

Definition A.20. Consider a square matrix $A = (a_{ij})$. Consider also the graph with n vertices given by set $X = \{x_1, \dots, x_n\}$. There is a path joining a vertex i to vertex j if $a_{ji} \neq 0$. Conversely, to a directed graph we associate a square matrix $n \times n$, where $a_{ij} = 1$ if there is an arc that connects i and j and $a_{ij} = 0$, otherwise.

We can characterize an irreducible matrix elegantly by the properties of the associated directed graph.

Theorem A.13. A matrix A is irreducible if and only if its graph $\mathcal{G}(A)$ is strongly connected.

Proof : The necessary condition.

Suppose the matrix A is irreducible. Let i be a vertex, we define I as the set of vertices different from i that can be reached from i . In the set I for all vertices $j \neq i$ there exists an arc joining j to i . The set I is non empty. In effect, if we consider the set, J , Indeed, and if we consider the set J , the complementary of the singleton $\{i\}$, as A is irreducible there exists $k \notin J$ and $j \in J$ such that $a_{jk} \neq 0$. But given the definition of J , this means that there exists $j \neq i$ such that $a_{ji} \neq 0$. In other words there is an arc from i to j .

Suppose by contradiction that $I \neq \{1, \dots, n\}$. By irreducibility of A , there exists $j \in I$ and $k \notin I$ such that $a_{kj} \neq 0$. That is there is an arc from i to k . But as $j \in I$, it is accessible from i , so k is accessible from i . This is a contradiction.

Sufficient condition.

Assume by contradiction that the graph is strongly connected and the associated matrix A is reducible. Then there exists a subset of indices I , such that if J is its complement, then $a_{ji} = 0$ for all $i \in I$ and $j \in J$. We choose an index $i \in I$ and an index $j \in J$. This is always possible since I is a proper subset. Then there exists an arc joining i to j . Let there the set of indices $\{k_1, \dots, k_p\}$ be such that the following coefficients are non-zero

$$a_{j,k_1}; a_{k_1,k_2}, \dots, a_{k_p,i}.$$

From the assumptions on i and j , it follows that $a_{k_p,i} \neq 0$, $k_p \notin J$ or $k_p \in J$. But if $k_p \in I$ the same reasoning applied on a_{k_p,k_p} prove that $k_{p-1} \in I$. reasoning by induction shows that $j \in I$ which is a contradiction.

Proposition 4. A is an irreducible Metzler matrix then each trajectory of the system $\dot{x} = Ax$ are and will remain on one face. Precisely, A is an irreducible matrix, if and only if for any $t > 0$, we have $e^{tA} \gg 0$

Proof

We assume that it is not true. Then there exists a $t > 0$ such that $e^{tA} \in \partial \mathcal{R}_+^{n^2}$. But for all s such that $0 \leq s \leq t$ we have $e^{(t-s)A} e^{sA}$. The matrix $e^{(t-s)A}$ is positive because (A is Metzler) and it is invertible. consequently if $e^{sA} \gg 0$ this implies $e^{tA} \gg 0$. We now show that e^{sA} is not strongly positive for all $0 \leq s \leq t$. There exists $x > 0$ belonging to the boundary of the orthant, such that $e^{tA}x$ belongs to the boundary of the orthant. From the previous proposition, there is an index i such that $\langle e_i | e^{tA}x \rangle = 0$ and $\langle e_i | A e^{tA}x \rangle > 0$.

The function $\varphi(s) = \langle e_i | e^{sA}x \rangle$ is positive for all s . It vanishes at $t = s$. This is a minimum, thus the derivative vanishes at t . Now

$$\dot{\varphi}(t) = \langle e_i | e^{tA}x \rangle > 0.$$

This is a contradiction. Conversely, if A is reducible, then A can be expressed as :

$$P^T A P = \begin{bmatrix} A_1 & A_2 \\ 0 & A_4 \end{bmatrix},$$

or as

$$P^T e^{tA} P = \begin{bmatrix} e^{tA_1} & M(t) \\ 0 & e^{tA_4} \end{bmatrix}$$

The matrix e^{tA} is reducible and positive, and there exists a vector $x > 0$ on the boundary such that $e^{tA}x$ is on the boundary of the orthant. This means that we cannot verify that $e^{tA} \gg 0$.

A.4.4 Regular Splitting of a Metzler matrix

We present and prove a result of Varga [152], which is applied in the formula for calculating the basic reproduction number, $\mathcal{R}_0$.

Definition A.21. Let a Metzler matrix A be invertible. A regular decomposition is a decomposition of the form

$$A = F + V$$

where $F \geq 0$ and V is a Metzler stable matrix.

The following theorem is proved in [152], but given its importance in this thesis, we also present the proof here.

Theorem A.14. Let A be an invertible Metzler matrix. For a regular decomposition of A of the form $A = F + V$, the following two assertions are equivalent.

1. A is a stable matrix
2. $\rho(-F V^{-1}) < 1$

Proof

Suppose A is a Metzler matrix, then from theorem A.10, we have $-A^{-1} \geq 0$.

The matrix $V = A - F$ and A are invertible so we can write

$$-F V^{-1} = -F (A - F)^{-1} = -F A^{-1} (I - F A^{-1})^{-1}$$

Let $G = -F V^{-1}$ be a positive matrix. To get the spectral radius using the Perron-Frobenius theorem, it is sufficient to restrict ourselves to the positive eigenvectors.

So let $v > 0$ be an eigenvector of G corresponding to an eigenvalue $\lambda \geq 0$, so that $Gv = \lambda v$. Now,

$$-F V^{-1}v = G (I - G)^{-1}v = \frac{\lambda}{1 - \lambda} v$$

The matrix $-F V^{-1}$ is positive. Conversely, let $\mu \geq 0$ be an eigenvalue corresponding to an eigenvector $v > 0$. Then $G (I - G)^{-1}v = \mu v$. The matrix G and $(I - G)^{-1}$ commute, and we deduce that $Gv = \mu (I + G)v$. This necessarily implies that $\mu \neq 1$ and v is an eigenvector of G corresponding to the eigenvalue $\frac{\mu}{1+\mu}$.

The function defined by $x \rightarrow \frac{x}{1+x}$ in $[0, 1] \in \mathbb{R}^+$ is a bijection between the eigenvalues of $G = -F A^{-1}$ and those of $-F V^{-1}$. This is a monotonic function. Consequently we have

$$\rho(-F V^{-1}) = \frac{\rho(G)}{1 + \rho(G)} < 1.$$

Conversely, assume $\rho(-F V^{-1}) < 1$. Then the matrix $I - F V^{-1}$ is invertible and in addition a Metzler matrix. Since $\rho(-F V^{-1}) < 1$, we have $\alpha(-I - F V^{-1}) < 1$. This is a Metzler stable matrix. Its inverse is positive and therefore

$$-A^{-1} = (-I - F V^{-1})^{-1} V^{-1} \geq 0.$$

We see from theorem A.10 that A is a Metzler stable matrix, and this completes the proof.

A.5 Calculation of Basic Reproduction Number

To calculate the basic reproduction number we demonstrate here the method developed by van den Driessche and Watmough [149]. In the formula, $\mathcal{R}_0$ is defined as the spectral radius of Next Generation Operator. Determination of the operation involves distribution of the populations into two compartments, the compartment of infected (latent, infectious ...) and the compartment of uninfected individuals.

This technique was developed first by Diekmann and Heesterbeek in [32] and later improved by van den Driessche and Watmough [149] for finite dimensional systems.

Consider an epidemiological model with no classes or with homogeneous compartments. The vector x represents the state of the system and x_j is the number (or concentration) of individuals in compartment j . The compartments are sorted in such a way that the first k compartments are free of infection (susceptibles) while the others are the infected compartments (latents, infected.).

Set the vector $x = x_j$, $j = 1, \dots, n$ where x_j is the number of individuals in compartment j .

Let $\mathcal{F}_j(x)$ be the rate of appearance of new infections in compartment i , $\mathcal{V}_i^+(x)$ be the rate of transfer of individuals into compartment i by all other means, and $\mathcal{V}_j^-(x)$, the rate of transfer of individuals out of compartment j .

The dynamics of the compartment is defined by

$$\dot{x}_j = \mathcal{F}_j(x) + \mathcal{V}_j^+(x) - \mathcal{V}_j^-(x)$$

It is assumed that each function is continuously differentiable at least twice in each variable. If we put $\mathcal{V}_j(x) = \mathcal{V}_j^+(x) - \mathcal{V}_j^-(x)$ the previous system becomes

$$\dot{x}_j = \mathcal{F}_j(x) + \mathcal{V}_j(x),$$

if x_0 is the disease free state of the system, and the infected compartments are empty. This equilibrium is called the Disease Free Equilibrium, ie $j > k$, $(x_0)_j = 0$. For biological reasons we have the following hypotheses :

1. $x \geq 0$, $\mathcal{F}_j(x) \geq 0$, $\mathcal{V}_j^+(x) \geq 0$, $\mathcal{V}_j^-(x) \geq 0$. Since each function represents a directed transfer of individuals, they are all non-negative. If a compartment is empty, then there can be no transfer of individuals out of the compartment by death, infection, nor any other means. Therefore, the following also holds
2. If $x_i = 0$ then $\mathcal{V}_j^-(x) = 0$. In particular, we if set $X_s = \{x \geq 0; x_j = 0, i = 1, \dots, n\}$ and if $x \in X_s$, then $\mathcal{V}_j^-(x) = 0$. In other words, there can be no transfer from an empty compartment.

3. If $j \leq k$ then $\mathcal{F}_j(x) = 0$. That is, there is no immigration of infectives from the uninfected compartment.
4. If x_0 is the disease free state then $\mathcal{F}_j(x_0) = 0$ and for $j \geq k$, $\mathcal{V}_j^+(x_0) = 0$. If the population remains near the DFE ie. if there is no disease, then introduction of a few infected individual will not result in an epidemic.

We now estimate the average number of new infections resulting from introduction of a typical individual near the DFE. To determine the fate of a typical infective individual introduced into the population, we consider the dynamics of the linearized system with reinfection blocked. That is the system

$$\dot{x} = D\mathcal{V}(x_0)(x - x_0) = D\mathcal{V}^+(x_0)(x - x_0) - D\mathcal{V}^-(x_0)(x - x_0).$$

The following result is the precise structure of the linearised system $DX(x_0)$ near the disease free equilibrium x_0 .

Lemma 2. If x_0 is a DFE, then the matrices $D\mathcal{F}(x_0)$ and $D\mathcal{V}(x_0)$ are decomposed in blocks

$$D\mathcal{F}(x_0) = \begin{bmatrix} 0 & 0 \\ 0 & F \end{bmatrix}, \quad D\mathcal{V}(x_0) = \begin{bmatrix} J_1 & J_2 \\ 0 & V \end{bmatrix}$$

$F \geq 0$ and V is a Metzler stable matrix.

The matrix FV^{-1} is called the Second Generation Matrix. A complete proof of this theorem is done in [149].

Definition A.22 (Basic Reproduction Number, $\mathcal{R}_0$). The basic reproduction number, $\mathcal{R}_0$ is the spectral radius of the second generation matrix, namely

$$\mathcal{R}_0 = \rho(-FV^{-1})$$

The following interpretation is given of the matrix $-FV^{-1}$. Consider an infected individual introduced into compartment $k > m$ to a population without disease. The entry (i, k) of the matrix $-V^{-1}$ is the average length of time this individual spends in compartment j during its lifetime, assuming that the population remains near the DFE and re-infection is blocked. The (i, j) entry of F is the rate at which infected individuals in compartment j produce new infections in compartment i . Hence, the (i, k) entry of the product FV^{-1} is the expected number of new infections in compartment i produced by the infected individual originally introduced into compartment k .

BIOLOGY OF MOSQUITO AND MALARIA PARASITE

In this appendix, we describe the history and biology of malaria and the malaria transmitting mosquito. The word, malaria, is derived from the Italian phrase, ‘mal-aria’, meaning bad air.

Malaria infection is caused by a protozoan parasite of the genus *Plasmodium*, four species of which infect human beings with the most common being *Plasmodium vivax* and the most deadly being *Plasmodium falciparum*. The *Anopheles* mosquito serves as *Plasmodium*’s delivery system, or vector. Only female mosquitoes can transmit it since males do not take blood meals.

B.0.1 The parasites life cycle.

The malaria parasite exhibits a complex life cycle involving the mosquito and a human host. Infection begins when the female *Anopheles* mosquito bites an infected human and injects saliva with sporozoites into the human blood stream. Sporozoites are carried by the circulatory system to the liver and invade a variety of liver cells. The sporozoites replicate to produce merozoites which are released into the bloodstream. Merozoites invade erythrocytes and enlarge to produce trophozoites. The trophozoites subdivide and mature to become schizonts, which are released following a rupture of the infected erythrocyte. Invasion of erythrocytes reinitiates another round of the blood-stage replicative cycle, usually referred to as the asexual cycle. Sometimes, instead of the asexual replicative cycle, the parasite can differentiate into sexual forms known as macro- or microgametocytes. Ingestion of gametocytes by the mosquito vector induces gametogenesis (i.e., the production of gametes). Microgametes then fertilize the macrogamete to produce a zygote which develops into a motile ookinete. The ookinete will penetrate the gut wall cells and develop into an oocyst. The oocyst matures and releases the sporozoites which migrate to, and invade, the salivary glands. This completes the life cycle of the parasite cycle. The incubation period (time from infection to development of the disease) is usually about 10 to 15 days.

B.0.2 Mosquitoes Life cycle

Anopheles have four stages in their life cycle : egg, larva, pupa, and adult. The first three stages last 5-14 days depending on the species and the ambient temperature. The adult females lives for about 1-2 weeks in nature.

Eggs : An anopheles mosquito lays between 50-200 eggs per oviposition directly on water. The eggs have floats on either side to help them float till they hatch within 2-3 days, in ambient temperature.

Larvae : Mosquito larvae have a well-developed head with mouth brushes used for feeding, a large thorax, and a segmented abdomen. They breathe through spiracles located on the abdomen and therefore must come to the surface frequently. Larvae spend most of their time feeding on algae, bacteria, and other microorganisms in the surface microlayer. They dive below the surface only when disturbed. They develop through 4 stages, or instars, after which they metamorphose into pupae. At the end of each instar, the larvae molt, shedding their exoskeleton, or skin, to allow for further growth. The larvae prefer clean, unpolluted water and many species prefer habitats with vegetation. Some breed in open, sun-lit pools while others are found only in shaded breeding sites in forests.

Pupae : The pupae is comma-shaped when viewed from the side. The head and thorax are merged into a cephalothorax with the abdomen curving around underneath. They usually come to the surface frequently to breathe, through a pair of respiratory trumpets on the thorax. After a few days the pupa splits and the adult mosquito emerges. Once larvae emerge to become adults, the rate at which they feed on man is dependent upon the ambient temperature.

Adults : Anopheles mosquitoes can be distinguished from other mosquitoes by the palps, which are as long as the proboscis, and by the presence of discrete blocks of black and white scales on the wings. Adult Anopheles can also be identified by their typical resting position : they rest with their abdomens sticking up in the air rather than parallel to the surface on which they are resting. The duration from egg to adult varies considerably among species and is strongly influenced by ambient temperature. Mosquitoes can develop from egg to adult in as little as 5 days but usually take 10-14 days in tropical conditions.

Some Anopheles mosquitoes prefer to bite humans and others prefer animals ; they are usually referred to as anthropophilic and zoophilic respectively. The anthropophilic are more likely to spread malaria because they bite humans more often. Most Anopheles are active at dusk and dawn (crepuscular) while others are nocturnal (active mostly at night). Some anopheles prefer to feed/rest indoors (endophagic / endophilic), while others feed/rest outdoors (exophagic / exophilic). When feeding, the female mosquito first injects some saliva into the host to prevent clotting, allowing the malaria parasite to enter the human host.

Some Anopheles species are poor vectors of malaria, because the parasites do not develop well (or at all) within them. These refractory strains have an immune response that encapsulates and kills the parasites after they have invaded the mosquito's stomach wall. It is hoped that some day, genetically modified mosquitoes that are refractory to malaria can replace wild mosquitoes, thereby limiting or eliminating malaria transmission.

At 17°C the female mosquitoes feed on humans every 4 days while at 25°C they take blood meals from humans every 2 days. Rainfall increases the breeding habitats for mosquitoes thus leading to increased population sizes and the rate of malaria transmission.

The rate of the parasite development in the female mosquito has an exponential relationship to temperature, hence a very small increase in external temperature will reduce the time it takes for the parasite to mature several fold. In Western Kenya a 0.5°C increase in temperature since the 1970's can explain the eight-fold increase in malaria cases (<http://www.ipsnews.net/news>).

BIBLIOGRAPHIE

- [1] Y. Afrane, B. Lawson, A. Githeko, and G. Yan, Effects of microclimatic changes due to land use and land cover on the duration of gonotrophic cycles of *Anopheles gambiae* giles (diptera : Culicidae) in western kenya highlands., *J Med Entomol*, 42 (1998).
- [2] Y. Afrane, B. Lawson, A. Githeko, and G. Yan, Effects of microclimatic changes caused by land use and land cover on duration of gonotrophic cycles of *Anopheles gambiae* (diptera : Culicidae) in western kenya highlands., *J Med Entomol*, 42 (2005).
- [3] H. Alles, K. Mendis, and R. Carter, Malaria mortality rates in south asia and in africa : implications for malaria control, *Parasitology Today*, 14 (1998).
- [4] R. Anderson and R.M. May, *Infectious Diseases of Human Dynamics and Control*, Oxford Science Publications, 1991.
- [5] J. Angulo, C. Takiguti, C. Pederneiras, and P. Megale, Identification of pattern and process in the spread of a contagious disease, *Soc. Sci. Med*, 13D (1979), p. 183.
- [6] J. Arino, Diseases in metapopulations, in *Modeling and Dynamics of Infectious Diseases*, Z. Ma, Y. Zhou, and J. Wu, eds., vol. 11 of *Series in Contemporary Applied Mathematics*, World Scientific, 2009, pp. 65–123. Also CDM Preprint Series report 2008-04.
- [7] J. Arino, J. Davis, D. Hartley, R. Jordan, J. Miller, and P. van den Driessche, A multispecies epidemic model with spatial dynamics, *Math. Med. Biol.*, 22 (2005), p. 129.
- [8] J. Arino, R. Jordan, and P. van den Driessche, Quarantine in a multi-species epidemic model with spatial dynamics, *Mathematical Biosciences*, 206 (2007).
- [9] J. Arino and P. van den Driessche, A multi-city epidemic model, *Math. Pop. Stud.*, 10 (2003).
- [10] J. Arudo, J. E. Gimnig, F. O. ter Kuile, S. P. Kachur, L. Slutsker, M. S. Kolczak, W. A. Hawley, A. S. S. Orago, B. L. Nahlen, and P. A. Phillips-Howard, Comparison of government statistics and demographic surveillance to monitor mortality in children less than five years old in rural western kenya., *American Journal of Tropical Medicine and Hygiene*, 68 (2003), pp. 30–37.
- [11] P. Auger, E. Kouokam, G. Sallet, M. Tchuenté, and B. Tsanou, The Ross-Macdonald model in a patchy environment, *Mathematical Biosciences*, 216 (2008), pp. 123–131.
- [12] P. Auger, E. Kouokam, G. Sallet, M. Tchuenté, and B. Tsanou, The Ross-Macdonald model in a patchy environment, *Math. Biosci.*, 216 (2008), pp. 123–131.

- [13] N. Bailey, *The Biomathematics of Malaria.*, Springer, 2010.
- [14] N. J. T. Bailey, *The mathematical theory of infectious diseases and its application*, Macmillan Publishers, 1975.
- [15] M. Balls, R. Bodker, C. Thomas, W. Kisinza, H. Msangeni, and S. Lindsay, Effect of topography on the risk of malaria infection in the usambara mountains, tanzania., *Trans R Soc Trop Med Hyg*, 98 (2004).
- [16] M. Bayoh and S. Lindsay, Effect of temperature on the development of the aquatic stages of *anopheles gambiae sensu stricto* (diptera : Culicidae)., *Bull Entomol Res* 2003, 93(5) :375-381., 93(5) (2003), pp. 375–381.
- [17] N. Becker, The use of epidemic model., *Biometrics*, 35 (1978).
- [18] A. Berman and R. Plemmons., *Nonnegative Matrices in the Mathematical Sciences*, Academic Press, 1979.
- [19] A. Berman and R. J. Plemmons, *Nonnegative matrices in the mathematical sciences*, vol. 9 of *Classics in Applied Mathematics*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1994. Revised reprint of the 1979 original.
- [20] B.M. Bloom, *Tuberculosis : Pathogenesis, protection and control.*, ASM Press, 1994.
- [21] Bony and Jean-Michel, Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés, *Ann. Inst. Fourier (Grenoble)*, 19 (1969), pp. 277–304 xii.
- [22] T. Bousema and C. Drakeley, Epidemiology and infectivity of *plasmodium falciparum* and *plasmodium vivax* gametocytes in relation to malaria control and elimination., *Clin. Microbiol.*, 24(2) (2011), p. 77.
- [23] G. Butler and P. Waltman, Persistence in dynamical systems., *Proc. Amer. Math. Soc.*, 96 (1986).
- [24] L. M. Cai and X. Z. Li, Global analysis of a vector-host epidemic model with nonlinear incidences, *Applied Mathematics and Computation*, 217 (2010).
- [25] V. Capasso and G. Serio, A generalisation of the kermack-mckendrick deterministic epidemic model, *Math. Biosci*, 42 (1978).
- [26] N. Chitnis, J. M. Hyman, and J. M. Cushing, Determining important parameters in the spread of malaria through the sensitivity analysis of a mathematical model., *Bull. Math. Biology*, 70 (2008), pp. 1272–1296.
- [27] N. R. Chitnis, *Using Mathematical models in controlling the spread of malaria*, PhD thesis, University of Arizona, 2005.
- [28] S. Connor, M. Qinoes, M. Thomson, M. Jawara, J. Todd, and B. Greenwood, Movement of *anopheles gambiae s.l.* vectors between villages, *Soc. Sci Med*, 43 (1996).
- [29] C. Cosner and al., The effects of human movement on the persistence of vector- borne diseases., *J. Theor. Bioll*, 258 (2009).
- [30] F. Coutinho, M. Burattinia, L. Lopeza, and E. Massad, Threshold conditions for a non-autonomous epidemic system describing the population dynamics of dengue, *Bulletin of Mathematical Biology*, 68 (2006).
- [31] B. Daniel, Essai d’une nouvelle analyse de la mortalité causée par la petite vérole et des avantages de l’inoculation pour la prévenir., *Mem. Math. Phys. Acad. Roy. Sci.*, (1760), pp. 1–45.

- [32] O. Diekmann and J. Heesterbeek, Mathematical epidemiology of infectious diseases in Mathematical and Computational Biology, Wiley Series, 2000.
- [33] O. Diekmann, J. A. P. Heesterbeek, and J. A. J. Metz, On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations, *J. Math. Biol.*, 28 (1990), pp. 365–382.
- [34] K. Dietz, Epidemics and rumours : A survey., *J. Roy. Statist. Soc. Ser. A*, 130 (1967).
- [35] ———, Overall population patterns in the transmission cycle of infectious disease agents, no. 87, Springer, Berlin, 1982.
- [36] K. Dietz, L. Molineaux, and A. Thomas, A malaria model tested in the african savannah, *Bulletin of the World Health Organization*, 50 (1974).
- [37] K. Dietz and D. Schenzle, Mathematical models for infectious disease statistics, in *A Celebration of Statistics.*, Springer-Verlag, 1985.
- [38] C. Drakeley, J. Bousema, N. Akim, K. Teelen, W. Roeffen, A. Lensen, M. Bolmer, W. Eling, and R. Sauerwei, Transmission-reducing immunity is inversely related to age in plasmodium falciparum gametocyte carriers., *Parasite immunology*, 28(5) (2006).
- [39] A. Ducrot, B. Some, S. Sirima, and P. Zong, A mathematical model for malaria involving differential susceptibility, exposedness and infectivity of human host, *Journal of Biological Dynamics*, (2008).
- [40] D.W.Kelly and C. E. Thompson, Epidemiology and optimal foraging : modelling the ideal free distribution of insect vectors., *Parasitology*, 120 (2000), pp. 319–327.
- [41] L. Esteva and C. Vargas, A model for dengue disease with variable human population, *J. Math. Biol.*, 38 (1999).
- [42] A. Fall, A. Iggdr, G. Sallet, and J. Tewa, Epidemiological models and lyapunov exponents, *Math.Model.Nat.Phenom.*, 2 (2007), pp. 62–83.
- [43] M. Fiedler, *Special Matrices and their Applications in Numerical Mathematics*, Martinus Nijhoff Publishers, 1986.
- [44] G. R. Fulford, M. G. Roberts, and J. A. P. Heesterbeek, The metapopulation dynamics of an infectious disease : tuberculosis in possums, *Theor Popul Biol*, 61 (2002).
- [45] P. Garnham, Malaria epidemics at exceptionally high altitude in kenya., *Br Med J*, 2 (1945).
- [46] G.D.Shanks, K. Biomndo, and S. et al., Changing patterns of clinical malaria since 1965 among a tea estate population located in the kenyan highlands, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 94 (2000).
- [47] P. Gething, T. V. Boeckel, D. Smith, C. Guerra, A. Patil, R. Snow, and S. Hay, Modelling the global constraints of temperature on transmission of plasmodium falciparum and p. vivax., *Parasit Vectors* 2011, 4 :92., 4 (2011), p. 92.
- [48] A. Githeko, D. Branding-Bennet, M. Beier, F. Atieli, M. Owaga, and F. Collins, The reservoir of plasmodium falciparum malaria in a holoendemic area of western kenya, *Action of Royal Society of Tropical Medicine and Hygiene*, 86 (1992), pp. 335–358.
- [49] A. Githeko and W. Ndegwa, Predicting malaria epidemics using climate data in kenyan highlands : a tool for decision makers., *Global Change and Human Health*, 2 (2001).
- [50] A. K. Githeko, J. M. Ayisi, P. K. Odada, F. K. Atieli, B. A. Ndenga, J. I. Githure, and G. Yan, Topography and malaria transmission heterogeneity in western kenya highlands : prospects for focal vector control, *Malar J.*, 5 (2006), p. 107.

- [51] B. Greenfell and J. Harwood, (meta)population dynamics of infectious diseases, Elsevier Science, 12 (1997).
- [52] S. Gupta., R. Snow, C.A.Donnely, K. Marsh, and Newbold, Immunity to non-cerebral severe malaria is acquired after one or two infections, *Nature Medicine*, 5 (1999), pp. 340–343.
- [53] R. Gurtler, L. Ceballos, P. O. Krasnowski, and L. L. R. Stariolo, Strong host-feeding preferences of the vector *triatoma infestans* modified by vector density : implications for the epidemiology of chagas disease., *PLoS Neglected Tropical Diseases* ., 3 (2009), p. 447.
- [54] H. L. Guyatt, K. Corlett, P.Timothy, Robinson, S. A. Ochola, and R. Snow, Malaria prevention in highland kenya : indoor residual house-spraying vs. insecticide-treated bednets., *Tropical Medicine and International Health*, 43 (2002).
- [55] J. Hale, *Ordinary Differential Equations*, New York, 1969.
- [56] J. Hale, *Ordinary differential equations*, Krieger, 1980.
- [57] S. I. Hay, A. M. Noor, M. Simba, M. Busolo, H. L. Guyatt, S. A. Ochola, and R. W. Snow, Clinical epidemiology of malaria in the highlands of western kenya, *Emerg. Infect. Dis.*, 8(6) (2002).
- [58] S. I. Hay, D. J. Rogers, S. E. Randolph, D. I. Stern, J. Cox, G. D. Shanks, and R. W. Snow, Hot topic or hot air? climate change and malaria resurgence in east african highlands., *RENDS in Parasitology Vol.18 No.12 December 2002*, 18(12) (2002).
- [59] J. Heesterbeek and J. Metz, The saturating contact rate in marriage and epidemic models, *J. Math. Biol*, 31 (1993).
- [60] H. Hethcote, Qualitative analysis of communicable disease models, *Mathematical Biosciences*, 28 (1976).
- [61] —, *A thousand and one epidemic models*, Springer-Verlag, 1994.
- [62] H. Hethcote and H. Thieme, Stability of endemic equilibrium in epidemic models with subpopulations, *Math. Biosci*, 75 (1985), pp. 205–227.
- [63] H. W. Hethcote, The mathematics of infectious diseases, *SIAM Rev*, 42(4) (2004).
- [64] Y. Himeidan, E. Malik, and E. Adam, Epidemiological and seasonal pattern of malaria in irrigated areas of eastern sudan., *Am J Infect Dis*, 1 (2005).
- [65] Y. Himeidan, G. Zhou, L. Yakob, Y. Afrane, S. Munga, H. Atieli, El-Rayah, A. Githeko, and G. Yan, Habitat stability and occurrences of malaria vector larvae in western kenya highlands., *Am J Infect Dis*, 8 (2009), p. 234.
- [66] M. Hirsch, Systems of differential equations that are competitive or cooperative i : limit sets, *SIAM J. Math. Anal.*, 13 (1982), pp. 167–179.
- [67] M. Hirsch, Systems of differential equations that are competitive or cooperative i : limit sets, *SIAM J Math Anal*, 13(2) (1982), p. 167.
- [68] M. Hirsch, The dynamical system approach to differential equations, *Bull. AMS*, 11 (1984), pp. 1–64.
- [69] —, Systems of differential equations that are competitive or cooperative ii : convergence almost everywhere, *SIAM J. Math. Anal.*, 16 (1985), pp. 423–439.
- [70] —, Systems of differential equations that are competitive or cooperative iii : Competing species, *Nonlinearity*, 1 (1988), pp. 51–71.

- [71] ———, Systems of differential equations that are competitive or cooperative. iv : Structural stability in three-dimensional systems, *SIAM J. Appl. Math.*, 21 (1990), pp. 1125–1234.
- [72] M. W. Hirsch and H. L. Smith, Monotone dynamical systems, in *Handbook of differential equations : ordinary differential equations. Vol. II*, Elsevier B. V., Amsterdam, 2005, pp. 239–357.
- [73] ———, Monotone dynamical systems, in *Handbook of differential equations : ordinary differential equations. Vol. II*, Elsevier B. V., Amsterdam, 2005, pp. 239–357.
- [74] C. Holling, The functional response of predators to prey density and its role in mimicry and population regulations., *emoirs of the Entomological Society of Canada*, 45 (1965), pp. 3–60.
- [75] P. Hviid, Natural acquired immunity to plasmodium falciparum malaria in africa, *Acta Tropica*, 95 (2005).
- [76] J. M. Hyman and J. Li, Differential susceptibility and infectivity epidemic models, *Mathematical Biosciences and Engineering*, 3 (2006), pp. 89–100.
- [77] IRIN, Kenya : Climate change and malaria in nairobi, 2012.
- [78] J. A. Jacquez and C. P. Simon, Qualitative theory of compartmental systems, *SIAM Rev.*, 35 (1993), pp. 43–79.
- [79] S. James, Report on a visit to kenya and uganda to advise on anti-malarial measures., tech. rep., London, 1929.
- [80] V. Jansen and A. Lloyd, Local stability analysis of spatially homogeneous solutions of multi-patch systems, *J. Math. Biol.*, 41 (2004).
- [81] C. Jost, O. Arino, and R. Arditi, About deterministic extinction in a ratio-dependent predator-prey model., *Bulletin of Mathematical Biology*, 61 (1999), pp. 19–32.
- [82] D. Kelly and C. Thompson., Epidemiology and optimal foraging : modelling the ideal free distribution of insect vectors., *Parasitology*, 120 (2000), pp. 319–327.
- [83] W. Kermack and A. G. McKendrick, Contributions to the mathematical theory of epidemics ii. 1927., *Proc R Soc London*, 115 (1927), pp. 700–721.
- [84] ———, Contributions to the mathematical theory of epidemics i. 1927., *Proc R Soc London*, 138 (1932), pp. 55–83.
- [85] ———, Contributions to the mathematical theory of epidemics iii. ., *Proc R Soc London*, 141 (1933).
- [86] K.Ghosh, C. J., and T. P.K., Immunity boosted by low exposure to infection in ansirs model, *Eiological Modelling*, 87 (1996).
- [87] G. F. Killeen, B. G. J. Knols, and W. Gu, Taking malaria transmission out of the bottle : Implications of mosquito dispersal for vector control interventions, *THE LANCET Infectious Diseases*, 3 (2003).
- [88] M. Kirby and Lindsay, Effect of temperature and inter-specific competition on the development and survival of anopheles gambiae sensu stricto and an. arabiensis larvae., *Acta Tropica* 2009, 109(2) :118-123., 109(2) (2009), pp. 118–123.
- [89] M. J. Klowden, The endogenous regulation of mosquito reproductive behavior, *Experientia*, 46 (1990), pp. 660–670.
- [90] K.Marsh, L. Otoo, and R. Hayes, Antibodies to blood stages antigens of plasmodium falciparum in rural gambians and their relationship to protection against infection, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 83 (1989).

- [91] M. Kretzschmar, J. Wallinga, M. Gail, K. Krickeberg, J. Samet, A. Tsatis, and W. Wong, *Mathematical Models in Infectious Disease Epidemiology*, Springer, 2010.
- [92] C. Kribs-Zaleta, To switch or taper off : the dynamics of saturation., *Mathematical Biosciences* 192 : 137-152., 192 (2004), pp. 137–152.
- [93] C. Kribs-Zaleta, Vector consumption and contact process saturation in sylvatic transmission of t. cruzi, *Mathematical Population Studies*, 13 (2006).
- [94] C. Kribs-Zaleta, Sharpness of saturation in harvesting and predation., *Mathematical Biosciences and Engineering* 6 : 719-742., 6 (2009), pp. 719–742.
- [95] C. Kribs-Zaleta, Estimating contact process saturation in sylvatic transmission of trypanosoma cruzi in the united states, *PLoS Neglected Tropical Diseases*, 4 (2010), p. 656.
- [96] A. Lajmanovich and J. Yorke, A deterministic model for gonorrhea in a nonhomogeneous population., *Math. Biosci.*, 28 (1976.), pp. 221–236.
- [97] J. LaSalle, Stability theory for ordinary differential equations. stability theory for ordinary differential equations., *J. Differ. Equations*, 41 (1968), pp. 57–65.
- [98] J. Lasalle., *The Stability of Dynamical Systems in : Society for Industrial and Applied Mathematics.*, Regional conference series in Applied Mathematics, 1976.
- [99] J. LaSalle, Stability of nonautonomous systems., *Nonlinear Anal., Theory, Methods Appl.*, 1(1) (1976.), pp. 83–91.
- [100] J. P. LaSalle, *The stability of dynamical systems*, Society for Industrial and Applied Mathematics, Philadelphia, Pa., 1976. With an appendix : “Limiting equations and stability of nonautonomous ordinary differential equations” by Z. Artstein, Regional Conference Series in Applied Mathematics.
- [101] P. Leenheer and H. Smith, Virus dynamics : A global analysis, *SIAM J Appl.Math*, 63 No. 4 (2003), pp. 1313–1327.
- [102] Levins, *Evolution in Changing Environments.*, Princeton University Press, 1968.
- [103] M. Li and J. Mouldowney, Global stability for the seir model in epidemiology, *Mathematical Biosciences*, 125 (1995).
- [104] S. Lindsay and W. Martens, Malaria in the african highlands : past, present and future., *B World Health Organ*, 76 (1998).
- [105] F. Lopez-Antunano, Epidemiology and control of malaria and other arthropod-borne diseases., *Mem Inst Oswaldo Cruz*, 87(Suppl 3) (1992), pp. 105–114.
- [106] G. Macdonald, *The Epidemiology and control of malaria.*, Oxford University Press, 1957.
- [107] M. A. Malakooti, K. Biomndo, and Shanks, Reemergence of epidemic malaria in the highlands of western kenya, *Emerging Infectious Diseases*, 4 (1998), p. 4.
- [108] M. Markus and H. Minc., *A survey of matrix theory and matrix inequalities.*, Dover, 1 (1992).
- [109] K. Marsh, Malaria : A neglected disease., *Parasitology*, 104 (1992).
- [110] A. Matson, The history of malaria in nandi, *East Afr Med J.*, 34 (1957).
- [111] C. McCluskey and P. van den Driessche, Global analysis of tuberculosis models, *Journal of Differential Equations*, 16 (2004).
- [112] C. C. McCluskey, Global stability for an sir epidemic model with delay and nonlinear incidence, *Nonlinear Anal. RWA*, 11 (2010).

- [113] McMichael, Githeko, McCarthy, O. Canziani, N. Leary, D. Dokken, and K.S.White, Human Health. In *Climate Change 2001, Impacts, Adaptations and Vulnerability of Climate Change. Contribution of working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change.*, Cambridge University Press, 2001.
- [114] A. L. Menach, F. E. McKenzie, A. Flahault, and D. L. Smith, The unexpected importance of mosquito oviposition behaviour for malaria : non-productive larval habitats can be sources for malaria transmission, *Malaria Journal*, 4 (2005), p. 23.
- [115] M. Janet, C. Mbogo, H. Mwambi, M. D. Wilson, G. Ojwang, J. M. Mwangangi, J. G. Nzovu, J. G. G. Yan, and J. C. Beier, Estimating dispersal and survival of anopheles gambiae and anopheles funestus along the kenyan coast by using mark-release-recapture methods, *J Med Entomol*, 44(6) (2007).
- [116] J. Mouldowney, Compound matrices and ordinary differential equations, *Rocky Mountain Journal of Mathematics*, 20 (1990).
- [117] W. R. Mukabana, W. Takken, R. Coe, and B. G. J. Knols, Host-specific cues cause differential attractiveness of kenyan men to the african malaria vector anopheles gambiae., *Malar J*, 1 (2002), p. 17.
- [118] B. Ndenga, A. Githeko, E. Omukunda, G. Munyekenye, H. Atieli, P. Wamai, C. Mbogo, N. Minakawa, G. Zhou, and G. yan, Population dynamics of malaria vectors in western kenya highlands., *J. Med. Entomology*, 43 (2006), pp. 200–206.
- [119] G. Ngwa and W. Shu, A mathematical model for endemic malaria with variable human and mosquito populations, *Math. Comput. Model*, 32 (2000).
- [120] G. A. Ngwa, On the population dynamics of the malaria vector, *Bulletin of Mathematical Biology*, 68 (2006).
- [121] S. Omar, I. Adagu, D. Gump, N. Ndaru, and D. Warhust, Plasmodium falciparum in kenya : high prevalence of drug resistance- associated polymorphisms in hospital admissions with severe malaria in an epidemic area., *Ann Trop Med Parasitol*, 95 (2001).
- [122] M. Pascual, J. Ahumada, L. Chaves, X. Rodo, and M. Bouma, Malaria resurgence in the east african highlands : Temperature trends revisited., *Proc Nat Acad Sci USA*, 103 (2006).
- [123] J. Patz, M. Hulme, C. Rosenzweig, T. Mitchell, R. Goldberg, A. Githeko, S. Lele, A. McMichael, and D. L. Sueur, Regional warming and malaria resurgence., *Nature*, 420 (2002).
- [124] R. E. Paul, S. Bonnet, C. Boudin, T. Tchuinkam, and V. Robert, Age-structured gametocyte allocation links immunity to epidemiology in malaria parasites, *Malaria Journal*, 6 (2007), p. 123.
- [125] L. Petersen and J. Roehrig, West nile virus : a reemerging world health organization, a multicentre collaboration to investigate the cause of severe acute respiratory syndrome, *Lancet*, 361 (2003), p. 1730.
- [126] P. Pongsumpun and I. Tang., Transmission of dengue haemorrhagic fever in an age structured population, *Mathematical and Computer Modeling*, 37 (2003), pp. 949–961.
- [127] M. Quincampoix, Differential inclusions and target problems, *SIAM J. Control Optim.*, 30 (1992), pp. 324–335.
- [128] R. Ross, *The prevention of Malaria*, Springer-Verlag, 1911.
- [129] Ruan, W. Wang, and S. Levin., The effect of global travel on the spread of sars, *Math. Biosci. Eng.*, 3(1) (2006).

- [130] J. Saldana, Continuous time formulation of reaction-diffusion processes on heterogeneous metapopulations., *Phys. Rev. E*, 78 (2008).
- [131] ———, Modelling the spread of infectious diseases in complex metapopulations., *Math. Mod. Nat. Phen.*, 5 (2010), p. 6.
- [132] G. Sallet and J. Kamgang, Global asymptotic stability for the disease free equilibrium for epidemiological models., *C. R. Math. Acad. Sci*, 341 (2005), p. 433.
- [133] L. Sattenspiel and K. Dietz., A structured epidemic model incorporating geographic mobility among regions, *Math. Bio. Sci.*, 128(1)(2) (1995).
- [134] M. Service., Mosquito (diptera : Culicidae) dispersal the long and short of it., *J Med Entomology*, 34 (1997), pp. 579–588.
- [135] G. Shanks, S. Hay, D. Stern, and et al, Meteorologic influences on plasmodium falciparum malaria in the highland tea estates of Kericho, western Kenya., *Emerging Infectious Diseases*, 12 (2002).
- [136] S.I.Hay, M. Simba, M. Busolo, A. Noor, H. Guyatt, S. Ochola, and R. Snow, Defining and detecting malaria epidemics in the highlands of western Kenya, *Emerg. Infect. Dis.*, 8 (2002).
- [137] D. L. Smith, C. A. Guerra, R. W. Snow, and S. I. Hay, Standardizing estimates of the plasmodium falciparum parasite rate., *Malar J*, 6 (2007), p. 131.
- [138] H. L. Smith, Monotone dynamical systems : an introduction to the theory of competitive and cooperative systems., *Mathematical Surveys and Monographs*. 41. Providence, RI : American Mathematical Society (AMS). x, 174 p. , 1995.
- [139] S. Staedke, E. Nottingham, J. C. Kamya, M. Rosenthal, and P. Dorsey, Short report : proximity to mosquito breeding sites as a risk factor for clinical malaria episodes in an urban cohort of Ugandan children., *Am J Trop Med Hyg*, 69 (2003), pp. 244–246.
- [140] H. Thieme, Global asymptotic stability in epidemic models, in *Equadiff82*, Springer-Verlag, 1993.
- [141] ———, Persistence under relaxed point-dissipativity (with application to an endemic model), *SIAM J Math Anal.*, 24 (1993), pp. 407–435.
- [142] ———, *Mathematics in Population Biology*, Princeton University Press, 2003.
- [143] H. R. Thieme, Spectral bound and reproduction number for infinite population structure and time-heterogeneity, *SIAM J. Appl. Math.*, 70 (2009), pp. 188–211.
- [144] J. Trape, G. Pison, M. Preziosi, C. Enel, L. Desgrees, V. Delaunay, B. Samb, E. Lagarde, J. Molez, and F. Simondon, Impact of chloroquine resistance on malaria mortality., *C. R. Acad. Sci III*, 321 (1998), pp. 689–697.
- [145] J. Tumwiine, J. Mugisha, and L. Luboobi, A mathematical model for the dynamics of malaria in a human host and mosquito vector with temporary immunity, *Applied Mathematics and Computation*, 189 (2007).
- [146] J. Tumwiine, J. Y. T. Mugisha, and L. S. Luboobi., On oscillatory pattern of malaria dynamics in a population with temporary immunity, *Computational and Mathematical Methods in Medicine*, 8 (2007).
- [147] UNICEF, Kenya statistics, tech. rep., United Nations, 2010.
- [148] P. van den Driessche and J. Watmough, reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, *Math. Biosci.*, 180 (2002), pp. 29–48.

- [149] P. van den Driessche and J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, *Math. Biosci.*, 80 (2002), p. 29.
- [150] J. van Dillen, M. Custers, A. Wensink, B. Wouters, T. van Voorthuizen, W. Voorn, B. Khan, L. Muller, and C. Nevill, A comparison of amodiaquine and sulfadoxine-pyrimethamine as firstline treatment of falciparum malaria in kenya., *Trans R Soc Trop Med Hyg*, 93 (1999).
- [151] R. Varga, Factorisation and normalised iterative methods *Boundary Problems in Differential Equation*, University of Wisconsin Press, 1960.
- [152] ———, *Matrix Iterative Analysis.*, Prentice-Hall, 1962.
- [153] R. Varga, *Gershgorin and His Circles*, Apprentice Hall, Springer, 2004.
- [154] A. Vespagnani and V. Colizza, Epidemic modelling in metapopulation systems with heterogeneous pattern : theory and simulations, *J. Theor. Biol.*, 251 (2008), p. 450.
- [155] A. Vespagnani, V. Colizza, and A. Barrat, Modeling the world-wide spread of pandemic influenza, *PLoS Med.*, 4 (2007), p. 95.
- [156] A. Vespagnani, V. Colizza, and R. Pastor-Storras, Reaction-diffusion processes and metapopulation models in heterogeneous networks, *Nat. Phys.*, 3 (2007), p. 276.
- [157] M. Vidyasagar, Decomposition techniques for large-scale systems with nonadditive interactions : Stability and stabilizability., *IEEE Trans. Autom. Control*, 25 (1980), pp. 773–779.
- [158] M. Vidyasagar, Decomposition techniques for large scale sytems with nonadditive interactions : Stability and stabilization., *IEEE Trans. Autom. Control*, 25 (1980), p. 773.
- [159] W. Wang and G. Mulone, Threshold of disease transmission in a patchy environment, *J. Math. Anal. Appl.*, 285 (2002).
- [160] W. Wang and X. Zhao, An epidemic model in a patchy environment., *Math. Biosci.*, (2004), p. 190.
- [161] C. L. Wanjala, J. Waitumbi, G. Zhou, and A. K. Githeko, Identification of malaria transmission and epidemic hotspots in the western kenya highlands : its application to malaria epidemic prediction., *Parasites and Vectors*, 4 (2011), p. 81.
- [162] White, Modelling the impact of vector control interventions on anopheles gambiae population dynamics, *Parasites & Vectors*, 4 (2011), p. 153.
- [163] G. White, The anopheles gambiae complex and malaria transmission around kisumu, kenya., *Trans R Soc Trop Med Hyg*, 66 (1972).
- [164] WHO, Anonymous chemotherapy of malaria and resistance to antimalarials., tech. rep., World Health Organization, 1973.
- [165] ———, Online global pathogen, *Emerg. Inf. Dis.*, 7 (2001), p. 611.
- [166] R. Xu and Z. Ma, Global stability of a sir epidemic model with nonlinear incidence rate and time delay, *Nonlinear Anal. RWA*, 10 (2009).
- [167] J. A. Yorke, H. W. Hethcote, and A. Nold, Dynamics and control of the transmission of gonorrhea., *Sex Transm Dis*, 5 (1978), pp. 51–56.
- [168] J. Zhang and Z. Ma, Global dynamics of an seir epidemic model with saturating contact rate, *Nonlinear Anal. RWA*, 185 (2003).
- [169] G. Zhou, N. Minakawa, A. Githeko, and G. Yan, Association between climate variability and malaria epidemics in the east african highlands, *Proc Natl Acad Sci USA*, 24 (2004).